

A study of USG in the evaluation of renal colic patients at tertiary health care center

Digvijay Ghodake¹, Abhijit Nagapurkar^{2*}, Nikit Mehta³, Anil G Joshi⁴

¹Junior Resident, ²Assistant Professor, ³Associate Professor, ⁴Professor and HOD, Department of Radio diagnosis, B V M C and Hospital, Sangli, Maharashtra, INDIA.

Email: digvi4630@gmail.com

Abstract

Background: The emergency department investigation of a patient with acute flank pain associated with hematuria has traditionally included an excretory urogram to exclude or confirm the clinical suspicion of ureteric calculi. **Aims and Objectives:** To Study USG in the evaluation of Renal colic patients at tertiary health care center. **Methodology:** All patients presenting with symptoms typical of renal colic attending the medicine and surgical OPD in santhiram medical college and general hospital were included into study while Pregnant women excluded from the study. Plain X-ray KUB, Ultrasound and CT scan study were done in 104 patients presenting with symptoms typical of renal colic attending the medical and surgical OPD in santhiram medical college and general hospital, CT imaging was performed using Siemens somatom Emotion duo machine. The data was analyzed by T-test in statistical package for the social science (SPSS). **Result:** USG positive in 74 people-71.15%, As per site Kidney-positive in 36 people-97%, Ureter-positive in 26 people - 48.1%, Bladder-positive in 6 people-100%, Urethra -positive in 6 people-100%. CT is superior to ultrasound and radiograph in all areas ultrasound is more sensitive for detecting renal, bladder and urethra calculi radiograph is more sensitive for detecting ureter, bladder and urethra calculi when compared to renal calculi. CT imaging is superior to ultrasound (97%) and radiograph (55%) for detection of calculi in kidney. For detection of calculi in ureter CT is superior to radiograph (64%) and ultrasound (48%). CT and ultrasound are equally sensitive and superior to radiograph for detecting urethral calculi. **Conclusion:** It can be concluded from our study that CT was superior to USG and Radiograph for overall detection of renal calculi but USG was superior to radiograph and if the condition of hydro nephrosis is considered the USG is useful as CT found useful in the management of renal calculi. **Key Words:** Renal colic, Renal calculi, UTI (Urinary Tract Infections), Hydro nephrosis.

*Address for Correspondence:

Dr. Abhijit Nagapurkar, Assistant Professor, Department of Radio diagnosis, B V M C and Hospital, Sangli, Maharashtra, INDIA.

Email: digvi4630@gmail.com

Received Date: 06/10/2017 Revised Date: 10/11/2017 Accepted Date: 07/12/2017

DOI: <https://doi.org/10.26611/1004124>

Access this article online	
Quick Response Code:	Website: www.medpulse.in
	Accessed Date: 17 December 2017

INTRODUCTION

The emergency department investigation of a patient with acute flank pain associated with hematuria has traditionally included an excretory urogram to exclude or confirm the clinical suspicion of ureteric calculi.¹ This investigation, however, has a number of relative

contraindications, including significant renal insufficiency, dehydration, past reaction to contrast agents, and pregnancy.² The availability of nonionic contrast media has reduced the risk of reaction but has added significantly to the cost of investigation. Ultrasound is a safe, noninvasive imaging modality that is used to evaluate intraperitoneal and retroperitoneal structures successfully.³ It has also been used to study the urinary tract.⁴ Limited studies have been published concerning its role in the diagnosis of renal calculi.⁵⁻⁷

MATERIAL AND METHODS

All patients presenting with symptoms typical of renal colic attending the medicine and surgical OPD in santhiram medical college and general hospital were included into study while Pregnant women excluded from the study. Plain X-ray KUB, Ultrasound and CT scan study were done in 104 patients presenting with

symptoms typical of renal colic attending the medical and surgical OPD in santhiram medical college and general hospital Plain X-ray KUB was taken after bowel preparation and on empty stomach. Films were taken in 500 ma Siemens klinoscopy-H machine. Ultrasound imaging was performed in the department of radiodiagnosis santhiram medical college and general hospital, using real time ultrasound machine (Esaote-Mylab 50x vision) which is equipped with 2.6,3.5 and 5

MHZ and HP image point Hz which is equipped with 2 to 10 MHz and curvilinear probes. Patients were examined in empty stomach and full urinary bladder in the supine position. CT imaging was performed in the department of radiodiagnosis santhiram medical college and general hospital, using Siemens somatom Emotion duo machine. The data was analyzed by T-test in statistical package for the social science (SPSS).

RESULT

Table 1: Ultrasound Vs CT

		CT calculus					Total	%
		Kidney	Ureter	Bladder	Absent	Urethra		
USG Calculus	Kidney	33	3	0	0	0	36	97
	Ureter	1	25	0	0	0	26	48.1
	Bladder	0	0	6	0	0	6	100
	Absent	6	20	0	4	0	30	
	Urethra	0	0	0	0	6	6	100
Total		40	48	6	4	6	104	

(P<0.05)

In present study USG positive in 74 people-71.15%, As per site Kidney-positive in 36 people-97%, Ureter-positive in 26 people-48.1%, Bladder-positive in 6 people-100%, Urethra -positive in 6 people-100%.




Figure 1: Comparison between Radiograph, Ultrasound and CT

In present study CT is superior to ultrasound and radiograph in all areas ultrasound is more sensitive for detecting renal, bladder and urethra calculi radiograph is more sensitive for detecting ureter, bladder and urethra calculi when compared to renal calculi. In present study CT imaging is superior to ultrasound (97%) and radiograph (55%) for detection of calculi in kidney. For detection of calculi in ureter CT is superior to radiograph (64%) and ultrasound (48%). In present study Ultrasound has poor rate of detection when compared to radiograph and CT for ureteric calculi. In present study CT and ultrasound are equally sensitive and superior to radiograph for detecting urethral calculi.

DISCUSSION

The role of ultrasound to delineate renal structure is well recognized. Ultrasound can detect renal tract dilatation with a high degree of accuracy.⁹ Ultrasound can also detect renal calculi, although difficulties arise in the midureter where the calculus and associated hydroureter may be obscured by overlying bowel gas. Previous comparisons of ultrasound and excretory urograms have given conflicting results. Hill *et al*¹⁰ studied 61 patients referred for outpatient excretory urography. Forty-one of the 61 were subsequently found to have a calculus. Ultrasound detected 66% of the calculi, while the excretory urogram detected 85%. Ultrasound was able to detect calculi at the ureterovesicle junction better than the excretory urogram. It should be noted that these patients were not studied in emergency situations, and their

preparation for examination was optimal. None of their patients had surgery. Ultrasound has been found to be less accurate in detection of renal stones compared to CT scan.¹¹ However, unilateral hydronephrosis in addition to finding of stone on ultrasound increases its sensitivity to 81.3-82.4% in identifying nephrolithiasis.^{12,13} We included finding of unilateral hydronephrosis with or without nephrolithiasis as a criteria for an “abnormal ultrasound”. Use of ultrasound as an initial imaging study is also justified based on several studies which have shown no difference in patient management and outcomes between ultrasound and CT scan in a patient with suspected nephrolithiasis.¹⁴⁻¹⁷ Ultrasound was found to be 97% sensitive in predicting need for surgical intervention when it showed a stone and/or hydronephrosis in patients presenting with renal colic.¹⁴ In other studies, rate of urological intervention was significantly lower^{16,18} or no patients required admission within 30 days¹⁷ in those with normal results on ultrasound. In our study we have seen that USG positive in 74 people-71.15%, As per site Kidney-positive in 36 people-97%, Ureter-positive in 26 people -48.1%, Bladder-positive in 6 people-100%, Urethra -positive in 6 people-100% CT is superior to ultrasound and radiograph in all areas ultrasound is more sensitive for detecting renal, bladder and urethra calculi radiograph is more sensitive for detecting ureter, bladder and urethra calculi when compared to renal calculi. CT imaging is superior to ultrasound (97%) and radiograph (55%) for detection of calculi in kidney. For detection of calculi in ureter CT is superior to radiograph (64%) and ultrasound (48%). In present study Ultrasound has poor rate of detection when compared to radiograph and CT. CT and ultrasound are equally sensitive and superior to radiograph for detecting urethral calculi.

CONCLUSION

It can be concluded from our study that CT was superior to USG and Radiograph for overall detection of renal calculi but USG was superior to radiograph and if the condition of hydro nephrosis is considered the USG is useful as CT, and found useful in the management of renal calculi.

REFERENCES

1. Rous SN: A review of 171 consecutive patients with urinary lithiasis. *J Urol* 1981; 126: 376-379.
2. Anseil G, Tweedie MCK, West CR, et al: The current status of reactions to intravenous contrast media. *Invest Radiol* 1980;15(supplJ): 532-539.
3. Kuligowska E, Newman B, White SJ, et al: Interventional ultrasound in detection and treatment of renal inflammatory disease. *Radiology* 1983; 147:521-526.
4. Rosenfield AT, Taylor KJW, Dembner AG, et al: Ultrasound of renal sinus: New observations. *Am J Radiol* 1979; 133:441-448.
5. Edell S, Zegel H: Ultrasonic detection of renal calculi. *AJR* 1978; 130:261-263.
6. Pollack HM, Arger PM, Goldberg BB, et al: Ultrasonic detection of nonopaque renal calculi. *Radiology* 1978; 127:233-237.
7. Diamant MJ, Malekzadeh M: Ultrasound and the diagnosis of renal and ureteral calculi. *Pediatrics* 1986; 109:980-983.
8. Sackett DL, Haynes RB, Tugwell P: *Clinical Epidemiology: A Basic Science for Clinical Medicine*. Toronto, Little, Brown and Co, 1985, p 22-28.
9. Ellenbogen PH, Scheible FW, Talner LB, et al: Sensitivity of grey scale ultrasound in detecting urinary tract obstruction. *AJR* 1978; 130:731-733.
10. Hill MC, Rich JJ, Mardiat JG, et al: Sonography vs excretory urography in acute flank pain. *AJR* 1985; 144:1235-1238.
11. Erwin BC, Carroll BA, Sommer PG: Renal colic: The role of ultrasound in initial evaluation. *Radiology* 1984; 152:147-150.
12. Fowler KA, Locken JA, Duchesne JH, Williamson MR: US for detecting renal calculi with non-enhanced CT as a reference standard. *Radiology*. 2002; 222:109-113.
13. P Riddell J, Case A, Wopat R, Beckham S, Lucas M, McClung CD, et al. Sensitivity of emergency bedside ultrasound to detect hydronephrosis in patients with computed tomography-proven stones. *West J Emerg Med*. 2014; 15:96-100. doi:10.5811/westjem.2013.9.15874.
14. Kanno T, Kubota M, Sakamoto H, Nishiyama R, Okada T, Higashi Y, et al. Determining the efficacy of ultrasonography for the detection of ureteral stone. *Urology*. 2014; 84:533-537.
15. Taylor M, Woo MY, Pageau P, McInnes MD, Watterson J, Thompson J, et al. Ultrasonography for the prediction of urological surgical intervention in patients with renal colic. *Emerg Med J*. 2016;33:118-123. doi:10.1136/emmermed-2014-204524.
16. Edmonds ML, Yan JW, Sedran RJ, McLeod SL, Theakston KD. The utility of renal ultrasonography in the diagnosis of renal colic in emergency department patients. *CJEM*. 2010; 12:201-206.
17. Fields JM, Fischer JJ, Anderson KL, Mangili A, Panebianco NL, Dean AJ, et al. The ability of renal ultrasound and ureteral jet evaluation to predict 30-day outcomes in patients with suspected nephrolithiasis. *Am J Emerg Med*. 2015; 33:1402-1406. doi:10.1016/j.ajem.2015.07.014.
18. Yan JW, McLeod SL, Edmonds ML, Sedran RJ, Theakston KD. Normal renal sonogram identifies renal colic patients at low risk for urologic intervention: a prospective cohort study. *CJEM*. 2015; 17:38-45.

Source of Support: None Declared
Conflict of Interest: None Declared