

Efficiency of intramuscular tramadol injection as labour analgesic and labour accelerator in latent phase of labour and mode of delivery

Maheshwari Marisiddaiah

Assistant Professor, Department of Obstetrics and Gynaecology, Bharatha Ratna Dr B.R. Ambedkar Medical College and Hospital, Bengaluru, Karnataka, INDIA.

Email: dr.mahe.m@gmail.com

Abstract

To prove the efficiency of Intramuscular Tramadol injection as labour analgesic and labour accelerator in latent phase of labour and mode of delivery. This is a prospective study done during Oct 2014 to Sep 2015. 100 pregnant women who were booked in this hospital were taken for study. Out of 100 patients, 26% had mild pain relief, 38% had moderate pain relief and 36% had no pain relief. Mean duration of I stage of labour was 256.09 minutes and II stage was 25.54 minutes. Mode of delivery – Normal delivery was 88.1% and LSCS was 9.1%

Key Words: Tramadol Injection, Latent phase of Labour, Labour Analgesia, Normal delivery, cesarean delivery.

Address for Correspondence:

Dr. Maheshwari Marisiddaiah, Assistant Professor, Department of Obstetrics and Gynaecology, Bharatha Ratna Dr B.R. Ambedkar Medical College and Hospital, Bengaluru, Karnataka, INDIA.

Email: dr.mahe.m@gmail.com

Received Date: 15/01/2018 Revised Date: 21/02/2018 Accepted Date: 10/03/2018

DOI: <https://doi.org/10.26611/1012613>

Access this article online

Quick Response Code:



Website:
www.medpulse.in

Accessed Date:
07 April 2018

emotional and sensory experience. It is of two types. Acute and Chronic. In 1847, Ether was introduced as first obstetrical anaesthetic agent by James Young Simpson.² In first stage of labour, perception of pain begins with nociceptive stimuli from mechanical and chemo receptors in cervix and uterus.³ In later stages repeated contraction produces myocellular injury which releases histamine, *i* serotonin, bradykinin, substance P, potassium ions and acetylcholine which stimulates chemical nociceptors.⁴ Pain score is recorded by patients history. The pain score should be reliable, valid. Dolorimeter is used to measure pain.⁵

INTRODUCTION

Labour pain is due to physiological, psychological excitatory and inhibitory complex interactions. Normal labour is series of events which results in expulsion of products of conception at term via natural or spontaneous onset of singleton vertex presentation.¹ Painless labour is usually preferred by every mother and it has positive influence on the course of labour. Epidural analgesia has been popularly used in western countries but it has limitations in India due to lack of trained staff and monitoring facilities. Intramuscular Tramadol has been used because of its convenience of administration and good, faster administration. Pain is an unpleasant

MATERIAL AND METHODS

This study is a prospective observational study done in the Dept of Obstetrics and Gynaecology in Bharatha Ratna Dr B.R. Ambedkar Medical College and Hospital, Bengaluru, over a period of 1 year from Oct 2014 to Sep 2015. About 100 term gestation women who were in latent phase of labour admitted to labour ward taken into the study. Details of the each women, degree of pain relief, duration of each stage of labour and mode of delivery were assessed at the end of each delivery.

RESULTS

In this study, out of 100 women, mean age of patients belongs to 21-25 y (53%), Parity distribution about 58(58%) were primi and 42(42%) were multiparous. 26 % had mild pain relief, 38% had moderate pain relief and 36% had no pain relief. Period of gestation between 37-39weeks were 47(47%) and 40-42weeks were 53(53%), Mean duration of I stage was 256.09 minutes and II stage was 25.54 minutes. Mode of delivery – normal delivery was 88.1% and LSCS was 9.1%.

DISCUSSION

An ideal method for pain relief should combine safety for both mother and fetus. The present study was designed to evaluate efficacy of IM Tramadol in latent phase of labour. The patients were standardized and compared regarding degree of pain relief duration of labour, and mode of delivery. In this study mean age of patients belongs to 21-25 y (53%). The difference in age group was not significant (p- 0.799). In some other study done by Thakur Ratna *et al* (2004), *i* the mean age was 22 years. In this study parity distribution about 58(58%) were primi and 42(42%) were multiparous. Period of gestation between 37-39weeks were 47(47%) and 40-42weeks were 53(53%). The variation in gestational age was not significantly significant (p- 0.087). The mean period of gestational age was 39 weeks in the study done by Sudha Patil *et al* (2014). In this study, 26 % had mild pain relief, 38% had moderate pain relief and 36% had no pain relief. In the study done by Thakur Ratna *et al*(2012) 16% had mild pain relief, 55% has moderate pain relief and 16% had no pain relief. The variation was statistically significant (p< 0.0001) In this study, Mean duration of I stage was 256.09 minutes and II stage was 25.54 minutes. In the study done by Khooshide h m alishahriari (2009)

mean duration of I stage was 140 minutes and II stage was 25 minutes. The variation was statistically significant (p<0.0001). In this study, Mode of delivery – normal delivery was 88.1% and LSCS was 9.1%. in the study done by O Kuti *et al* (2008) normal delivery was 88%, *i* LSCS was 9.5% and instrumental delivery was 2.4% and was significant.

CONCLUSION

Intramuscular Tramaol injection is effective in pain relief and it also shortens the duration of first and second stage of labour. Injection Tramadol didn't cause any major side effects in either mother or fetus and also no major increase in LSCS.

ACKNOWLEDGMENT

Authors acknowledge the scholars whose articles are cited and included in references of this manuscript. The authors are also grateful to authors/editors/publishers of all those articles and journals from where the literature for this article has been reviewed and discussed

REFERENCES

1. Friedman E.A. Primigravid labour: A graphicostastistical analysis *Obstet Gynecol* 6:567, 1955.
2. Simpson W. The works of Sir J Y Simpson Edinburg, Scotland; Adam and Charles Black 1871.
3. Bonica JJ, i McDonald JS. The pain of child birth. In:JJ Bonica, iEditor, The management of pain, Lea and Febiger, Philadelphia 1990;2:1313-1343
4. Crawford JS. Principles and practice of obstetric analgesia and anaesthesia. Blackwell publishers, Oxford, 1985;5:221.
5. Javert CT and Hardy J D. Am J Obst and Gynaec 60:552, 1950.

Source of Support: None Declared

Conflict of Interest: None Declared