

A cross section study of road traffic accidents with respect to road users' type, offending vehicles and cause of death among autopsies conducted at tertiary hospital mortuary rims Raichur

Sharanabasavappa Karaddi¹, Suraj^{2*}, Sujit N Jadhav³, Ravishankar M G⁴

¹Associate Professor,^{2,4}Final year Postgraduate, Department of Forensic Medicine and Toxicology Raichur Institute of Medical Sciences Raichur, Karnataka Pin code -584102, INDIA.

³Assistant Professor, Department of Forensic Medicine and Toxicology, St Paul's Hospital Millennium Medical College, Addis Ababa, ETHIOPIA.

Email: drsujitkalgurji@gmail.com

Abstract

The present study was a cross section study conducted at tertiary hospital mortuary block of RIMS Hospital, Raichur during one-year period from 01-12-2014 to 30-11-2015. During the study period a total of 125 victims of RTA with various severities of injuries who died and autopsied at mortuary of RIMS Hospital, Raichur were studied. The purpose of the study was to know the pattern of road users and offending vehicles, period of survival of victims following the accident and to establish the relation between the type of skeletal injury and cause of death.

Key Word: RTA, Road users type, offending vehicle.

*Address for Correspondence:

Dr. Suraj, Final year Postgraduate, Department of Forensic Medicine and Toxicology Raichur Institute of Medical Sciences Raichur, Karnataka Pin code -584102, INDIA.

Email: drsujitkalgurji@gmail.com

Received Date: 08/04/2019 Revised Date: 10/05/2019 Accepted Date: 02/07/2019

DOI: <https://doi.org/10.26611/10181112>

Access this article online

Quick Response Code:



Website:

www.medpulse.in

Accessed Date:
16 July 2019

INTRODUCTION

The Global status report on road safety by the World Health Organization (WHO) published in 2013, is the second broad assessment of the status of road safety in 182 countries, building on the first Global status report on road safety published in 2009, the latter of which used data drawn from a standardized survey conducted in

2008. Large regional disparities exist in the death rates of road traffic accidents, the risk being highest in the African region (24.2 per 100,000 population), and the lowest in Europe (10.3 per 100,000). Young adults (between 15 and 44 years) account for almost 60 per cent of all traffic deaths, and half of the world's road traffic deaths occur among pedestrians (22 per cent), cyclists (5 per cent) and motorcyclists (23 per cent). Proportion of deaths among different road user types show, however, considerable inter- and intraregional variation.^{1,2} Road Traffic Accident (RTA) is any vehicular accident occurring on the roadway (i.e. originating on, terminating on, or involving a vehicle partially on the roadway). This includes collision of an automobile with a pedestrian, another automobile or with a non-automobile on the roadway or fall from a moving vehicle causing injuries or death of involved individuals. Pedestrian is any person involved in a vehicular accident who was not at the time of the accident riding in or on a motor vehicle, or animal

How to cite this article: Sharanabasavappa Karaddi, Suraj, Sujit N Jadhav, Ravishankar M G. A cross section study of road traffic accidents with respect to road users' type, offending vehicles and cause of death among autopsies conducted at tertiary hospital mortuary rims Raichur. *MedPulse International Journal of Forensic Medicine*. July 2019; 11(1): 06-13.
<https://www.medpulse.in/Forensic%20Medicine/>

drawn vehicle, or on a pedal cycle. Pedal cycle is any land transport vehicle operated solely by pedals (includes bicycle and tricycle). Pedal cyclist is any person riding on a pedal cycle or in a sidecar or trailer attached to such a vehicle. Motorcycle means two-wheeled motor vehicle, inclusive of any detachable sidecar having an extra wheel, attached to the motor vehicle. Motorcyclist is any person riding on a motorcycle or in a sidecar attached to such a vehicle. Light motor vehicle means any public or private transport vehicle (passenger or goods) of unladen weight not exceeding 7,500 kgs e.g. auto rickshaw, car, jeep, taxi etc. Medium motor vehicle means any public or private transport vehicle (passenger or goods) of unladen weight more than 7,500 kgs but not exceeding 12,000 kgs. E.g. Metador, Tempo, Van, Tractor, Cruiser, Bolero etc. Heavy motor vehicle means any public or private transport vehicle (passenger or goods) of unladen weight more than 12,000 kgs. E.g. Bus, truck, etc. Animal drawn vehicle is any non-motor vehicle driven by any animal. E.g. Tanga, Bullock cart etc

MATERIALS AND METHOD

The present study was a Prospective study. The study material comprised of victims of RTA who died in spot or died in RIMS Hospital and autopsied in the autopsy block (mortuary) of RIMS Hospital, Raichur during the 1-year period from 01 December 2014 to 30 November 2015. Ethical clearance for the present study was obtained from the Institutional Ethical Committee RIMS, Raichur. In the present study information regarding the bio-data of the deceased and various characters regarding the circumstances of the accident like type of victim / road user, type of offending vehicle, time of death were gathered from all possible sources like police records, hospital records and also by direct interrogation with investigating officer, eye witnesses (if available), relatives and friends of the deceased accompanying dead bodies. In each case, a thorough external and internal examination was done for fractures and other injuries and opinion as to the cause of death was made after the examination. The data thus obtained was recorded in the predesigned and pretested Proforma, and analysed.

RESULTS

During the present study a total of 425 medico legal autopsies were conducted out of which 125 cases were of RTA with skeletal injury. All the victims of RTA had fracture of one or the other bone. Deaths due to RTAs accounted for (26.88 %) of the total medico legal autopsies conducted. (Table 1) Out of 125 cases of fatal RTA, maximum 60 cases (48%) were motorcycle riders, followed by pedestrians 26 cases (20.8%) minimum were front seat passengers of four wheelers 6 cases (4.8%).

(Table 2) Heavy motor vehicle topped the list of offending vehicles 49 cases (39.2%) followed by MMV 36 cases (28.8%), LMV 13 cases (10.4%), Motor cycle 10 cases (8%), Others 10 case (8%), Unknown (Hit and Run) 7 cases (5.6%). (Table 3) Most common vehicle causing motorcycle rider casualty was Medium Motor Vehicle (MMV-Cruiser, Innova, Bolero) 20 cases (33.33%), followed by Heavy Motor Vehicle (HMV-Bus, Lorry, Cranes) 16 cases (26.66%), Light motor vehicle (LMV-Auto, Tum Tum etc) 10 cases (16.66%), Others (hitting stationery objects like pole, wall etc) 7 cases (11.66%) , Motorcycle with motorcycle 5 cases (8.33%) , Unknown (Hit and Run) cases 2 (3.33%). (Table 4) Most common offending vehicle causing pillion rider casualty was HMV 3 cases (30%) and unknown (hit and run) 3cases (30%). Motorcycle was the offending vehicle in 2 cases (20%), Light motor vehicle 1 case (10%), Medium motor vehicle 1 case (10%). (Table 5) Most common offending vehicle causing pedestrian casualty was HMV 12 cases (45.15%) followed by MMV 8 cases (30.76%). Motorcycle was the offending vehicle in 3 cases (11.53%), Unknown vehicles was the offending vehicle in 3 cases (11.53%). (Table 6) Most common offending vehicle causing Driver casualty was HMV 9 cases (64.28%) followed by LMV 2 cases (14.28%). MMV was the offending vehicle in 2 cases (14.28%), Unknown vehicles was the offending vehicle in 1 cases (7.14%). (Table 7) In my study a total of 15 passengers died due to fatal RTA, out of which 6 were FSP and 9 were RSP. The most common offending vehicle was MMV (especially the Cruiser), in which 9 deaths were occurred followed by LMV in 5 cases and HMV 1 case. (Table 8)

In our study out of 125 cases of RTA, Spot death were observed in 43 cases (34.4%), 44 cases (35.2%) died on the way to hospital and in 38 cases (30.4%) died in the hospital. (Table 9) In our study, 44 victims (34.2%) died within 1 day (24 Hours) after the accident. The maximum number of deaths were observed between 1 day to 3 days (74 cases, 59.2%). 6 victims (4.8%) survived beyond 4 days but died within 1 week. Only 1 (0.8%) victim survived beyond 1 week but died within 2 weeks. (Table 10) In the present study the cranial injury was observed in 45 cases, out of which in 39 cases (86.66%) the cause of death was attributed alone to cranial injuries where as in remaining 6 cases (13.33%) along with cranial injuries other fatal injuries were observed. Death due to multiple injuries were accounted for 42 cases (33.6%) where in injury to lungs, pelvic bone fractures, thigh and leg bone fractures, abdominal and pelvic visceral injuries were observed in varying combinations. Death due to haemorrhagic shock was associated with 37 cases (29.6%). One person died due to Septicaemia. (Table 11)

Table 1: Profile of Medico Legal Autopsies Conducted During The Study Period

Type of Case	Number	Percentage
RTA with Skeletal injury	125	26.88
Poisoning	90	19.35
Burns	76	16.34
Drowning	17	3.7
Railways	47	10.1
Snake bite	11	2.3
Electrocution	10	2.1
Assault	8	1.7
Hanging	19	4.1
Natural Death	40	8.63
Others	22	4.8
Total	465	100

Table 2: Accident Victims / Road User Type

Accident Victim Type	No of Case	%
Rider	60	48
Pillion	10	8
Pedestrian	26	20.8
Driver	14	11.2
Front Seat Passenger	6	4.8
Rear Seat Passenger	9	7.2
Total	125	100

Table 3: offending vehicles

Type of vehicle	No of cases	%
Motorcycle	10	8
LMV	13	10.4
MMV	36	28.8
HMV	49	39.2
Unknown	7	5.6
Others	10	8
Total	125	100

Table 4: vehicles causing motorcycle rider casualty

Offending vehicle	Number of cases	Percentage
Motor cycle	5	8.33
LMV	10	16.66
MMV	20	33.33
HMV	16	26.56
Unknown	2	3.22
Others	7	11.62
Total	60	100 %

Table 5: vehicles causing pillion rider casualty

Offending vehicle	Number of cases	Percentage
Motor cycle	2	20
LMV	1	10
MMV	1	10
HMV	3	30
Unknown	3	30
Total	10	100%

Table 6: vehicles causing pedestrian casualty

Offending vehicle	Number of cases	Percentage
Motor cycle	3	11.53
MMV	8	30.72
HMV	12	46.08
Unknown	3	11.53
Total	26	100 %

Table 7: vehicles causing driver casualty

Offending vehicle	Number of cases	Percentage
LMV	2	14.28
MMV	2	14.28
HMV	9	64.26
Unknown	1	7.14
Total	14	100%

Table 8: vehicles causing fsp/rsp casualty

Offending vehicle	Number of cases	Percentage
LMV	5	33.30
MMV	9	59.94
HMV	1	6.66
Total	15	100%

Table 9: place of death of victims

Place	No of Cases	%
On the spot	43	34.4
On the way to hospital	44	35.2
In the hospital	38	30.4
Total	125	100

Table 10: period of survival following accident

Site	No of cases	%
< 24 Hours	44	35.2
1 day-3 day	74	59.2
4 days- 7 days	6	4.8
10 days- 14 days	1	0.8
Total	125	100

Table 11: Relations Between Fracture And Cod

S. No	Cause of Death	No of cases	%
1	Cranial Cause alone (Head Injury)	45	36
2	Multiple Injuries	42	33.6
3	Haemorrhagic Shock	37	29.6
4	Septicaemia	1	0.8
	Total	125	100

DISCUSSION

Cause wise analysis of road accidents revealed that most of road accidents were due to over-speeding accounting for 36.8% of total accidents which caused 48,654 deaths and injuries to 1,81,582 persons. Dangerous/careless driving or overtaking caused 1,37,808 road accidents which resulted in 42,127 deaths and injuries to 1,38,533 persons during 2014. 3.2% of road accidents were due to poor weather condition. Driving under influence of

drug/alcohol contributed 1.6% of total road accidents which rendered 7,398 persons injured and 2,591 deaths in the country.³

Profile Of Medico Legal Autopsies Of Fatal Rta (Table No.1): During the period of present study, it was observed that deaths due to RTA accounted for 26.88 % of total medico legal autopsies conducted i.e. more than 25% of unnatural deaths were due to RTAs. This result is less when compared with results of studies conducted at,

Government Medical College, Jammu (48.92%)⁴; Government Medical College, Chandigarh (42.18%)⁵; AIIMS, New Delhi (35.5%)⁶; PGIMS, Rohtak (29.8%)⁷; MLN Medical College, Allahabad (42.29%)⁸; KMC, Manipal (36.50%)⁹; The results are **more** when compared with the results of studies conducted at RM College, Loni (24.1%)¹⁰; Office of Judicial Medical Officer, Colombo (22.6%)¹¹; P.D.U. Medical College and Associated Hospital, Rajkot (13.8%)¹². The difference in the number of RTA related deaths observed in different studies can be explained by the fact that RTA depends upon various epidemiological factors like geographical area, conditions prevailing in that region, category of road users, condition of road etc. Since Raichur is a very backward area High speed of driving by victims, bad road conditions with too many speed breakers with no proper signs, Improper enforcement by the traffic police might have contributed to this percentage of death.

Accident victim/road user type (table no.2): In the present study motorcycle riders formed the largest group of victims (48%). Maximum number of rider casualties can be explained by the fact that our country is on a progressive path and the population is now able to afford purchase two wheelers further more banks offering purchase of vehicles on EMI basis has made it easier than before. Each house hold is having minimum of two to three vehicle this increase in vehicles creates congestion on roads, rash speed driving by youngster all lead to increase of casualty in this group. Our study is similar to the results of study done at KMC Manipal, Karnataka where motor cyclists were involved in maximum number of cases (39.6%)⁹. However In the following studies pedestrians were more than motorcycle riders, Government Medical College, Jammu (46.17%)⁴; Government Medical College, Chandigarh (42.14%)¹³; Greater Lusaka Zambia (46.34%)¹⁴; Maulana Azad Medical college, New Delhi (39.1%)¹⁵. In the study conducted at Brisbane Queensland maximum number of victims were car drivers (32.98%)¹⁶ which is contrary to our result.

Offending Vehicles (Table No.3): In our study the most common offending vehicle was heavy motor vehicle (39.2%). Involvement of Heavy motor vehicle in accidents can be attributed to their high speed, presence of single space roads, allowing HMV into the congested centre of the city roads, fatigability, intoxication etc. Similar finding was also observed in other studies conducted at Government medical college, Jammu⁴; Government medical college, Chandigarh⁵; AIIMS, New Delhi⁶; PGIMS, Rohtak⁷; RM college, Loni¹⁰. During 2014, two wheelers have accounted for maximum fatal road accidents, contributing 26.4% of total road accidental deaths, followed by trucks/lorries (20.1%), cars

(12.1%) and buses (8.8%). 14.4% and 13.5% of deaths due to two wheel vehicles were reported in Tamil Nadu and Maharashtra respectively. 17.7% of deaths due to trucks/lorries and 15.5% of deaths due to accidents by cars were reported in Uttar Pradesh and Tamil Nadu respectively. Buses were responsible for 15.9% and 13.4% of fatal road accidents in Tamil Nadu and Uttar Pradesh respectively.³

Motor cycle rider vs offending vehicles (table no.4):

When we observe the table no.4, it is clear that the motor cycle rider causality was caused by all vehicles i.e Medium, Heavy and Low motor vehicles in various proportions. The congested roads, allowing heavy motor vehicles in the city limits, high speed and uncontrolled motor cycle riding, not following the traffic rules and avoiding helmets for various reasons might have increased the death rate in motor cycle riders in RTA. Study conducted by Naveen kumar et.al in Varanasi observed that 37% of death has occurred in two wheeler riders.¹⁷ Jan Mohammadi et.al in a study at Northern Iran observed that the death of Motor Cycle Riders was of 84.1% among total RTA.¹⁸ Michael Fitzharis, Rakhi Dandona conducted a study on 378 motorised two wheeled vehicles (MTV) users were enrolled in a study of whom 252 (66.7%) were riders.¹⁹

Pillion Rider Vs Offending Vehicles (Table No.5): Jan Mohammadi et.al in a study at Northern Iran observed that the death of pillion riders was of 15.7% among total RTA.¹⁸

Pedestrians vs offending vehicles (table no.6): Most common offending vehicle causing pedestrian casualty was HMV 12 cases (45.15%) followed by MMV 8 cases (30.76%). Motorcycle was the offending vehicle in 3 cases (11.53%), Unknown vehicles was the offending vehicle in 3 cases (11.53%). In an analysis of 101 pedestrian traffic deaths done at Czechoslovakia, higher occurrence of fracture of pelvis, femur and tibia was observed in accidents involving light motor vehicles and relatively more skull fracture, thoracic cage fracture and fracture of upper limb bones were seen in accidents involving medium and heavy motor vehicles.²⁰ Pedestrians' deaths in road accidents were reported as 17.5% and 12.5% in Maharashtra and Kerala respectively.³ Study conducted by Kyada HC et.al in Rajkot city observed that pedestrians were the commonest victims of RTA.²¹

drivers vs offending vehicles (table no.7): In the present study it was observed that occurrence of fracture of bones was higher in collision with heavy motor vehicles. This may be because of greater degree of force of impact due to large mass and more speed of heavy motor vehicles.

Front seat passengers/rear seat passengers vs offending vehicles (table no.8): In my study a total of 15

passengers died due to fatal RTA, out of which 6 were FSP and 9 were RSP. The most common offending vehicle was MMV (especially the Cruiser), in which 9 deaths were occurred followed by LMV in 5 cases and HMV 1 case. The unrestrained passenger move forward after collision and results in various skeletal injuries. The suddenness and force of impact will determine the severity of the injuries and survival of the victim. The violent impact of collision is more in rear seat occupants compared to front seat occupants since they are aware of what will happens next compared to rear seat occupants.

Place Of Death (Table No.9): In 54.8 % on the spot death cases heavy motor vehicles were the offenders. 84.1 % cases opted for treatment at tertiary centers. Injuries to chest/abdomen/pelvis proved to be fatal within 1 day. Deaths on the way to hospital were due to coma and hemorrhagic shock. These facts highlight the need for availability of prompt resuscitative measures and an approach of zero tolerance for traffic related hazards. This study finds that motor vehicle occupants suffered more severe injuries than the pedestrians. Whilst on the spot injuries requires preventive measures, a systematic approach needs to be incorporated to minimize deaths which occur on way to hospital and in the hospital. This study finds that on the way to hospital deaths were due to coma and hemorrhagic shock due to lack of prompt resuscitative measures in dealing with such emergencies. The success of the pre-hospital trauma management hinges on immediately making a triage and transport decision. Severely injured patients should be immediately transported to an appropriate hospital for definitive care using the "Load and go" philosophy, with all remaining care provided en route. Valuable pre-hospital care including a head to toe examination continuous monitoring, placement of subsequent intravenous access, and environmental control can be provided while the patient is being transported.¹⁷ In a study conducted by Singh YN et.al considering the place of death of the RTA victims, it is found that deaths in hospitals is highest (54.86%) which is more than the half of the total victims.²²

Period Of Survival Following Accident (Table No.10): In our study, maximum number of victims (35.2%) survived for a period of less than 24 hours. This finding emphasizes the fact that the first 24 hrs following the accident is the most crucial period, which decides the outcome in such cases. In the study conducted at GTB Hospital, Shahadra, New Delhi²³, 66% of victims died on the spot. Of the remaining victims, majority (45.44%) died within 24 hours, which is significantly higher than our result. In other study carried out at MLN Medical College, Allhabad⁸, maximum number of victims (40.21%) succumbed to the injuries sustained within 24

hrs after the accident. In another study conducted at Manipal, Karnataka⁹, highest number of victims (35.2%) died within 24 hours after the accident. The results of above studies are in general agreement with the results of our study.

Relation between fracture and cause of death (table no.11): In the present study the cranial injury was observed in 45 cases, out of which in 39 cases (86.66%) the cause of death was attributed alone to cranial injuries where as in remaining 6 cases (13.33%) along with cranial injuries other fatal injuries were observed. In our study 33.6 % of victims died due to multiple injuries. Deaths due to multiple injuries are injuries which are difficult to decide which was the most serious and mortal injury which lead to death of the victim, as in the crushed head with extrusion of brain or rupture of the aorta etc but in such cases it is quite acceptable to use the term 'multiple injuries', preferably listing several of the most lethal.²⁴ In our study 29.6% of victims died due to hemorrhagic shock. Fracture of multiple bones was present in all the victims who died due to hemorrhagic shock. This may be due to the sudden and severe loss of blood because of injury to neighbouring blood vessels by the fractured bones. (The blood loss is, one to two litres in fracture of multiple ribs, one to three litres in open fracture of thigh, half to two litres in closed fracture of thigh, half to one litre in closed fracture of lower leg, quarter to half in fracture dislocation of ankle and two to two and half litres in fracture of pelvis⁵). From the above results, it can be said that fractures in one or the other way, are associated with fatal outcome. In the study done at Birmingham accident Hospital, Birmingham²⁵, the skull was fractured in 72% of all head injury cases. According to a study done at GTB Hospital, Shahadra, New Delhi²³, during January -December 2003, head injury was the most common cause of death (60.96%) and fracture of skull was present in 80% of the victims who died due to head injury. In another study undertaken at Government Medical College, Jammu²⁶, head injury was responsible for majority of deaths (62.96%) and associated skull fracture was present in 93.18% cases. Fatal RTA were more prevalent on the secondary road system (47.97 per cent) and especially involved pedestrian and two wheeler vehicle users. Numbers of skeletal injuries (199) and internal organ injuries (202) exceeded the total number of victims (98) clearly indicating the multiplicity of injuries. The majority of RTA victims (n=46, 46.93%) died due to head injury. The study showed that most deaths in RTA, brought to a tertiary care rural hospital, took place either on the spot or within 24 hours of injury which is very alarming and highlights the need to take urgent steps to establish good pre-hospital care and provision of trauma

services at site.²⁷ Most of the diseased are riders (60%). Head and neck (56.22%) is the most common part to receive injuries in two wheeler accidents. Head injury (47.48%) is the most common cause of death. Most of the people are died between 6 to 24 hours (27.27%) after the accident.²⁸ An epidemiologic survey done by Ding SL *et al* of 2451 consecutive victims of traffic accidents, coming to or managed at, the emergency care department of a large Taipei metropolitan hospital in 1990. Major head injury fatalities occurred within one week of the collision.²⁹ A fracture of skull with associated brain injury is the most common cause of death but multiple injuries constitute a typical feature of motorcycle accidents. The use of crash helmets has reduced fatalities at low speeds but afford little or no protection at high speed. A full face helmet is better than an open face helmet.²⁶ In a study done by Banerjee KK *et al* out of 470 autopsies, 344 victims had traumatic fatality. The leading type of trauma was road traffic accident (RTA) (52%). Majority 258 of the victims had head injuries followed by thoracic injuries (169) and abdomen (125). Of the associated with trauma 24% of victims were died within the first three hours and majority 79% of the victims were reached the hospital and survived for a different interval.²³

CONCLUSION

Increased motorization, rampant encroachment of roads, nasty tendency of violating traffic rules and anarchic traffic systems have greatly contributed to rapid increase in RTAs. Population explosion is a catalysing factor for a number of accidents. Good conditioned roads, following traffic rules and awareness about RTA in the people will definitely bring down the incidence of RTA.

REFERENCES

- World Health Organization. Global status report on road safety: time for action. [Internet]. Geneva: World Health Organization; 2009 [cited 2016 Sep 12]. Available from: <http://public.eblib.com/choice/publicfullrecord.aspx?p=557632>
- WHO | Global status report on road safety 2013 [Internet]. [cited 2016 Sep 14]. Available from: http://www.who.int/violence_injury_prevention/road_safety_status/2013/en/
- Welcome to National Crime Records Bureau [Internet]. [cited 2016 Sep 12]. Available from: <http://ncrb.gov.in/>
- Road-traffic accidents--a demographic and topographic analysis. - PubMed - NCBI [Internet]. [cited 2016 Sep 15]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11506352>
- Sharma BR, Dasari H, Sharma V, Vij K. Dynamics of road traffic fatalities in Chandigarh – a surprise. Journal of Forensic Medicine and Toxicology 2002 Jan-Jun;19(1):25-29.
- Chandra J, Dogra TD, Dikshit PC. Pattern of Cranio-Intracranial injuries in Fatal Vehicular Accidents in Delhi, 1966-76, Med. Sci. Law 1979; 19(3) : 186-194.
- Journal of Indian Academy of Forensic Medicine [Internet]. [cited 2016 Sep 15]. Available from: <http://medind.nic.in/jal/t04/i1/jalt04i1c.shtml>
- Journal of Indian Academy of Forensic Medicine [Internet]. [cited 2016 Sep 15]. Available from: <http://medind.nic.in/jal/t05/i3/jalt05i3c.shtml>
- Palimar V, Arun M, Singh B, Mohanty MK. Victimologic study of road traffic fatalities. Medico Legal Update 2004 Jul-Sept; 4(#): 91-93.
- Kachre RV, Kachre VH, Asawa SS. Pattern of vehicular accidents in Pravara region: a rural region of Ahmadnagar district of Maharashtra. Journal of Forensic Medicine and Toxicology 2003 Jul-Dec; 20(2):29-32.
- Analysis of fatalities in road accidents [Internet]. [cited 2016 Sep 15]. Available from: <http://www.sciencedirect.com/science/article/pii/S0379073888902198>
- Pattern of Motorcyclist's Mortality in Mazandran Province, Northern Iran - Iranian Red Crescent Medical Journal - 2009 - Kowsar [Internet]. [cited 2016 Sep 15]. Available from: http://ircmj.com/?page=articleandarticle_id=179
- accident - definition of accident in English from the Oxford dictionary [Internet]. [cited 2016 Sep 14]. Available from: <http://www.oxforddictionaries.com/definition/english/accident>
- Patel NS. Traffic fatalities in Lusaka, Zambia. Med Sci Law 1979; 19(1):61-65. Epidemiological study of the victims of vehicular accidents in Delhi. - PubMed - NCBI [Internet]. [cited 2016 Sep 15]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/130401434>
- Epidemiological study of the victims of vehicular accidents in Delhi. - PubMed - NCBI [Internet]. [cited 2016 Sep 15]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/130401434>
- Injuries from road accidents. - PubMed - NCBI [Internet]. [cited 2016 Sep 15]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/13898956>
- Navin Kumar, Manoj Kumar, "Medicolegal Study of Fatal Road Traffic Accidents in Varanasi Region", International Journal of Science and Research (IJSR), www.ijsr.net, Volume 4 Issue 1, January 2015, 1492 - 1496 - See more at: http://www.ijsr.net/archive/v4i1/v4i1_03.php#sthash.buEQ7JyS.cpoJINqb.dpuf
- Pattern of Motorcyclist's Mortality in Mazandran Province, Northern Iran - Iranian Red Crescent Medical Journal - 2009 - Kowsar [Internet]. [cited 2016 Sep 15]. Available from: http://ircmj.com/?page=articleandarticle_id=179
- Crash characteristics and patterns of injury among hospitalized motorised two-wheeled vehicle users in urban India | BMC Public Health | Full Text [Internet]. [cited 2016 Sep 15]. Available from <http://bmcpublihealth.biomedcentral.com/articles/10.1186/1471-2458-9-11>
- Vorel F. Traffic injuries of pedestrians. The Journal of Czecho-Slovak Forensic Medicine 1993 [Cited 2006 Feb

- 06]; 38(4):32-6. Available from http://www.nemcb.cz/soudni/SLa/a1993_4.html-177k
21. Kyada HC, Mangal H, Momin SG, Vijapura MT, Bhuvu SD. Profile of road traffic accidents in Rajkot city. *J Ind Acad forensic Med* 2012;34(20):135-8
 22. Singh YN, Bairagi KK and Das KC. An epidemiological study of road traffic accident victims in medicolegal autopsies, *J Indian Acad Forensic Med*, 2005, 27(3); 166-169.
 23. Study of head injury victims in fatal road traffic accidents in Delhi. - PubMed NCBI [Internet]. [cited 2016 Sep 15]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10085617>
 24. Saukko P, Knight B. Knight's forensic pathology. 4th ed. London: Arnold; 2015:277-297
 25. Traffic-crash fatalities (1968-73): injury patterns and other factors. - PubMed NCBI [Internet]. [cited 2016 Sep 15]. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/840002>
 26. Parikh's Text Book of Medical Jurisprudence, Forensic Medicine and Toxicology. CBS Publishers and Distributors, 7th Edition, 2016 ; 327-328
 27. Pattern of injury in fatal road traffic accidents in a rural area of western Maharashtra, India. - PubMed - NCBI [Internet]. [cited 2016 Sep 12]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24133540>
 28. Jakkam Surender. Pattern of injuries in fatal road traffic accidents in Warangal area; *J Indian Acad Forensic Med*. Jan-March 2013 Vol 35. No 1
 29. [Head injuries in traffic accidents with emphasis on the comparisons between motorcycle-helmet users and non-users]. - PubMed - NCBI [Internet]. [cited 2016 Sep 15]. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7920094>.

Source of Support: None Declared
Conflict of Interest: None Declared

