

Detection of mycobacterium tuberculosis using sputum evaluator methods in a tertiary care medical college hospital

Shreeram Astic Deshpande^{1*}, Prasanna Gupta², Namrata Naithani³

^{1,2,3}Department of Microbiology, NIMS Medical College, Jaipur, Rajasthan, INDIA.

Email: shreeramad@rediffmail.com

Abstract

Background: Tuberculosis caused by *Mycobacterium tuberculosis* (or Koch's bacillus) even though curable and preventable it is still one of the leading infectious causes of morbidity and mortality especially in developing countries like India. Diagnostic methods available to detect tuberculosis are Direct approach by Microscopy, Culture, Biochemical test, Nuclear techniques. Sputum microscopy remains the mainstay of diagnosis of pulmonary tuberculosis but lacks sensitivity compared to culture. Pre-requisite for microscopic examination of sputum sample is prior decontamination. Rationale for the study: There are only a very few studies done in South India to report that Hypertonic Saline Sodium Hydroxide (HS-SH) method of decontamination is superior to gold standard NALC-NaOH method in India. So, we would like to evaluate Hypertonic Saline Sodium Hydroxide (HS-SH) method for decontamination of sputum samples for sputum microscopy. Main objective of our study is to evaluate the performance of the Novel Hypertonic Saline Sodium hydroxide (HS-SH) method with the gold standard NALC-NaOH for decontamination of sputum specimens to detect acid fast *Mycobacterium tuberculosis*. Institutional Human Ethical Committee clearance and Patient's informed written consent obtained. The Study was conducted as a prospective epidemiological study in Diagnostic Microbiology section of Central Service Laboratory for 6 months from November 11, 2015 to May 10, 2016. Total of 143 sputum samples obtained from both in-patients and out-patients of Karpagam Medical College Hospital from various departments like Medicine, Surgery, TB and RD and Orthopedics were subjected to decontamination by both HS-SH method and NALC-NAOH methods. Slides were then examined under oil immersion for acid fast bacilli by conventional Ziehl - Neelsen staining microscopic method. Results: Out of 143 microbiological sputum samples, 104 were found to be positive for acid fast bacilli by both Modified petroff's method and Hypertonic Saline Sodium Hydroxide method (HS-SH) of concentration and decontamination and 39 samples were negative by both these methods. Sensitivity, Specificity, Positive predictive value and Negative predictive value of the HS-SH method was found to be 100%. Conclusion:

Keywords: *Mycobacterium tuberculosis*, Sputum samples, for Hypertonic Saline - Sodium Hydroxide Method.

*Address for Correspondence:

Dr Shreeram Astic Deshpande, Department of Microbiology, NIMS Medical College, Jaipur, Rajasthan, INDIA.

Email: shreeramad@rediffmail.com

Received Date: 02/08/2020 Revised Date: 12/09/2020 Accepted Date: 14/10/2020

DOI: <https://doi.org/10.26611/10081631>

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). 

Access this article online

Quick Response Code:	Website: www.medpulse.in
	Accessed Date: 10 December 2020

INTRODUCTION

Tuberculosis caused by *Mycobacterium tuberculosis* (or Koch's bacillus) even though curable and preventable it is still one of the leading infectious causes of morbidity and mortality especially in developing countries like India. Global incidence of tuberculosis is 9.4 million cases annually, whereas India accounts for 2.0 million i.e., one-fifth of the Global incidence (2010). The Global prevalence rate is 3.8 million bacteriologically positive cases (2000) with death rate of 3, 25,000 per year due to TB. A total of 8.6 million patients newly developed tuberculosis in 2012 as estimated accounting for a

mortality of 1.3 million cases (Global Tuberculosis report, World Health Organization, 2013). In India the prevalence rate is 249 per lakh population (2009). The life time risk of a person developing Tuberculosis is 10% (10 out of 100). Diagnosis of Tuberculosis is by a combination of proper history taking, clinical examination, Microbiological evaluation by Sputum microscopy and Culture (Conventional or automated) which is the gold standard and also by correlation with roentgenologic examination. Resistance in Tuberculosis had become a global menace for treating physicians. Diagnostic methods available to detect tuberculosis are (ICMR Bulletin, 2002): Direct approaches are by Microscopy, Culture, Biochemical test, Nuclear techniques. In developing countries diagnosis of acid fast bacilli is performed by microscopic examination of Ziehl-Neelsen (ZN) stained sputum smear, because it is simple, inexpensive and it provides rapid results. But this technique has a low sensitivity 22-43% for single smear (Toman, 2004). And also up to 60% under optimal conditions (Asper *et al.*, 2003), when compared with cultures so that culture remains as gold standard to diagnose tuberculosis. The threshold for detection of acid fast bacilli in sputum samples under optimal condition is between 104-105 bacilli/ml, which can be obtained by decontamination and concentration techniques (IUATLD, 2005). Sputum microscopy remains the mainstay of diagnosis of pulmonary tuberculosis but lacks sensitivity compared to culture. Pre-requisite for microscopic examination of sputum sample is prior decontamination. Conventional methods for decontamination include Petroff's method which is routinely done, while N-Acetyl-L-Cysteine Sodium Hydroxide (NALC-NaOH) still remains a gold standard. N-Acetyl-L-Cysteine method has 4% NaOH and kills 60-70% of Mycobacteria, and the sensitivity for acid fast smear is 66.7%. Hypertonic Saline- Sodium Hydroxide (HS-SH) method has only 1.33% NaOH is used for decontamination (Ganoza *et al.*, 2008). Concentrated sputum techniques are much better than direct sputum microscopy and Modified Petroff's method is the widely used DC method but this method is too cumbersome to carry in field locations (Vasanthakumari *et al.*, 1988). Various decontamination methods used for decontamination of sputum samples for *Mycobacterium tuberculosis* are Bleach (6% NaOCl) method, Petroff's method, Modified Petroff's Sodium Hydroxide method, N-Acetyl-L-Cysteine- sodium Hydroxide method (NALC - NaOH), Hypertonic Saline Sodium Hydroxide(HS-SH) method, Zephiran- Trisodium Phosphate (Z-TSP) method, Oxalic Acid method, Sulphuric Acid method, Cetyl-Pyridinium Chloride - Sodium chloride (CPC) method and Phenyl Ammonium

sulphate (Ricaldi *et al.*, 2008; Singhal *et al.*, 2013). Relative Centrifugal Force for HS-SH method is high (4000rpm) and the sensitivity for acid fast bacilli smear is about 71.4% which is more than the N-acetyl-L-cysteine-Sodium hydroxide decontamination and concentration method. In hypertonic saline-NaOH method, hypertonic saline is used for mucolysis and Sodium hydroxide for decontamination (King *et al.*, 1997). Literature survey has shown HS-SH technique substantially increased the sensitivity of AFB smears compared with the routine direct smear for AFB and correlated well with the NALC-NaOH decontamination method for sputum culture in suspected Tuberculosis patients (King *et al.*, 1997; Ricaldi *et al.*, 2008). Hypertonic saline has been used successfully for the symptomatic treatment of cystic fibrosis. There are only a very few studies to report that Hypertonic Saline Sodium Hydroxide (HS-SH) method of decontamination is superior to gold standard NALC-NaOH method in India. So, we would like to evaluate Hypertonic Saline Sodium Hydroxide (HS-SH) method for decontamination of sputum samples for sputum microscopy. The main objectives of this study to evaluate the performance of the Novel Hypertonic Saline Sodium hydroxide (HS-SH) method with the gold standard NALC-NaOH for decontamination of sputum specimens to detect acid fast *Mycobacterium tuberculosis*.

MATERIALS AND METHODS

Institutional Human Ethical Committee clearance and Patient's informed written consent obtained. The Study was conducted as a prospective epidemiological study in Diagnostic Microbiology section of Central Service Laboratory of National Institute of Medical Sciences and Research (NIMS),Jaipur, for 6 months from December 2018 to April 2019. Total of 143 sputum samples obtained from both in-patients and out-patients of NIMS Medical College Hospital, Jaipur from various departments like Medicine, Surgery, TB and RD and Orthopedics were subjected to decontamination by both HS-SH method and NALC-NAOH methods. Slides were then examined under oil immersion for acid fast bacilli by conventional Ziehl-Neelsen staining microscopic method.

Inclusion Criteria

1. Patients with cough more than 2 weeks and/or blood stained sputum.
2. New sputum positive pulmonary tuberculosis patients before initiation of treatment.

Exclusion Criteria

1. Patient already under anti-tubercle drug treatment.

2. Inadequate sample volume (< 2ml).
3. Sample not representative of lower respiratory tract.
4. Patient denying participating in the study.

Decontamination Method

Procedure for Hypertonic Saline - Sodium Hydroxide Method (HS-SH) (Singhal *et al.*, 2013; Nora Morcillo *et al.*, 2008):

One ml portion of sputum was mixed with one ml 7 % (w/v) NaCl and one ml 4 % (w/v) NaOH in a sterile 15 ml centrifuge tube (BD Falcon) and homogenized for 15–20 s using a vortex mixer [Final concentrations (w/v) in 3 ml: 2.33 % NaCl, 1.33 % NaOH]. The tubes were then incubated at 37°C for 30 min. After incubation, the mixture was neutralized with sterile PBS (pH 6.8), bringing the total volume to 15 ml. The mixture was vortexed for 5 s and then centrifuged at 3400 g for 15 min at 15°C, using aerosol proof shields. The supernatant was discarded into a splash-proof container with a tuberculocidal solution. The procedure used has been described previously and is recommended by the Centers for Disease Control and Prevention and the WHO/IUATLD. one ml sputum was added to a 50 ml BD Falcon centrifuge tube with one ml of solution containing 0.5% (w/v) NALC, 2.67% (w/v) NaOH and 1.45% (w/v) sodium citrate and mixed well (final concentrations (w/v) in 2 ml: 0.25% NALC, 1.34% NaOH, 0.73% sodium citrate). The tubes were incubated at room temperature for 15 min. After incubation, the mixture was neutralized with PBS, bringing the total volume to 50 ml. The rest of the procedure was as described above. ZN sputum Smear preparation AFB sputum smear preparation done in Bio safety cabinet level II. One drop of each suspended pellet was used to prepare slides for AFB microscopy using the Ziehl–Neelsen stain. Each slide was coded, read blindly by a qualified technician and reported according to the National Tuberculosis Program and WHO/IUATLD standards (WHO, 1998). Smears were reported as follows: Grade 0 where no Acid Fast Bacilli (AFB) is observed in a total of 200 oil immersion fields, Scanty (Sc) where 1–9 AFB in 100 microscopic fields (few bacilli) is observed; 1+ with 10–99 AFB in 100 fields; 2+ with 1–10 AFB per field in at least 50 fields; 3+ with more than 10 AFB per field in at least 20 fields. Each slide will be coded, examined and graded according to the RNTCP guidelines.

RESULTS AND DISCUSSION

Out of 143 microbiological sputum samples, 104 were found to be positive for acid fast bacilli by both Modified petroff's method and Hypertonic Saline Sodium Hydroxide method (HS-SH) of concentration and decontamination and 39 samples were negative by both

these methods. Sensitivity, Specificity, Positive predictive value and Negative predictive value of the HS-SH method was found to be 100%.

Table.1 Comparison of N-Acetyl -L- Cysteine Sodium Hydroxide (NALC-NaOH) and Hypertonic Saline Sodium Hydroxide (HS-SH) methods: Decontamination methods NALC + - HSS

104	0
0	39

TOTAL 104 39 143

Decontamination methods for sputum microscopy for diagnosis of tuberculosis presently used are cumbersome, time consuming and expensive. Better methods are needed particularly developing countries. According to the study conducted by Christian Ganoza *et al.*, the sensitivity for acid-fast bacilli (AFB) smears had increased from 28.6% using the direct method to 71.4% Hypertonic Saline Sodium Hydroxide (HS–SH) method and 66.7% N-Acetyl-L-Cysteine (NALC-NaOH) using decontamination (DC) methods. In a study conducted in India, showed sensitivity of NALC-NaOH and HS-SH methods were 46% and 52% respectively (Sushil Kumar Chaudary, 2010). The proposed novel HS–SH DC method improved the sensitivity of AFB microscopy compared with a routine direct smear; its performance was comparable to that of the Modified Petroff's method for AFB smears, but it was methodologically simpler and less expensive, equally sensitive and can be adopted in National TB control programme especially in developing countries. Sensitivity, Specificity, Positive predictive value and Negative predictive value of the HS-SH method was found to be 100% which is comparable with the standard Modified petroff's method. Novel HS-SH method was not all cumbersome compared to the gold standard NALC-NaOH method. In conclusion, the HS-SH sputum decontamination method is equally sensitive, specific, cost-effective, feasible and less time consuming procedure compared to the gold standard NALC-NaOH method for decontamination of Mycobacterium tuberculosis from sputum samples and can be routinely used in all peripheral health centres because of the low cost, especially in developing countries.

REFERENCES

1. Global Tuberculosis report, World Health Organization, WHO Press, Geneva, Switzerland, 2013, pp.10-36.
2. Toman, K. 2004. Tuberculosis Case-Finding and Chemotherapy. Case Detection, treatment and monitoring – Questions and Answers, 2nd Edition Geneva WHO.
3. Asper, L., Mutsvangwa, J., Magwenzi, J., Chigara, N., Butterworth, A., Maso N, P. and Van der Stuyft, P. 2003. A comparison of direct microscopy, the concentration method and the Mycobacteria Growth Indicator Tube for the examination of sputum for acid fast bacilli. *Ent. J. Tuberc. Lung Dis.*, 7, 376-381.

4. IUATLD. 2005. Tuberculosis bacteriology-priorities and indications in high prevalence countries: position of the technical staff of the Tuberculosis Division of the International Union Against Tuberculosis and Lung Disease. *Int. J. Tuberculosis and Lung dis.*, 9, 355-361.
5. Ganoza, C.A., Ricaldi, J.N., Chauca, J., Rojas, G., Munayco, C., Agapito, J. *et al.*. 2008. Novel hypertonic saline- sodium hydroxide (HS-SH) method for decontamination and concentration of sputum samples for *Mycobacterium tuberculosis* microscopy and culture. *J. Med. Microbiol.*, 57: 1094-98.
6. Vasanthakumari, *et al.*. 1988. Concentrated sputum smear microscopy: A simple approach to better case detection in pulmonary tuberculosis. *Indian J. Tuberculosis*, 35: 80.
7. King, M., Dasgupta, B., Tomkiewicz, R.P. and Brown, N.E. 1997. Rheology of cystic fibrosis sputum after in vitro treatment with hypertonic saline alone and in combination with recombinant human deoxyribonuclease I. *Am. J. Respir. Crit. Care Med.*, 156, 173 –177. Ricaldi, J.N and Guerra, H. 2008. A simple and improved method for diagnosis of Tuberculosis using hypertonic saline and sodium hydroxide (HS-SH) to concentrate and decontaminate sputum. *Trop. Doct.*, 38: 97-99.
8. Singhal, R., M. Bhalla, S. Chakraborty, D. Beheram V.P. Myneedu. 2013. Phenol Ammonium sulfate microscopy method for diagnosis of pulmonary tuberculosis. *Int. J. Mycobacteriol.*, 2: 84-88.
9. Nora Morcillo, Belén Imperiale, Juan Carlos Palomino; New simple decontamination method improves microscopic detection and culture of mycobacteria in clinical practice. *Infection and Drug Resistance*, 1: 21-26.
10. Kent, P.T., and Kubica, G.P. 1985. 21-44.
11. Sushil Kumar Chaudary, Dept. of Microbiology, St. John's Medical college, Karnataka. 2010, "Comparison of Hypertonic Saline Sodium Hydroxide decontamination method with Modified Petroff's method for concentrating sputum samples"
12. ICMR Bulletin. 2002. Vol. 32, No.8, Part I: Techniques for diagnosis of T.B

Source of Support: None Declared
Conflict of Interest: None Declared

Policy for Articles with Open Access:

Authors who publish with MedPulse International Journal of Microbiology (Print ISSN: 2550-7648) (Online ISSN: 2636-4646) agree to the following terms: Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a Creative Commons Attribution License that allows others to share the work with an acknowledgement of the work's authorship and initial publication in this journal.

Authors are permitted and encouraged to post links to their work online (e.g., in institutional repositories or on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work.