

Role of MRI to assess the progress of spinal tuberculosis during the course of treatment

Saurabh Dwivedi^{1*}, Chandra Prakash Pal², Yogendra Kumar Mishra³, Tanu Mishra⁴

¹Resident, ²Head Of Department, ³Resident, Department of Orthopaedics, SN Medical College Agra, Uttar Pradesh, INDIA.

⁴Resident, Department of Radiology, RML Lucknow, Uttar Pradesh, INDIA.

Email: dwivedi7500@gmail.com

Abstract

Background: Spine is the commonest site of skeletal involvement in tuberculosis and constitutes more than 50% of all forms of osteo-articular tuberculosis. MRI is the most valuable non-invasive diagnostic and prognostic tool in the patients with spinal tuberculosis to promote early detection of disease and assess the response of treatment and change in regime of A.T.T. accordingly. **Objective:** Magnetic Resonance Imaging pattern of Spinal tuberculosis during course of treatment, to promote early detection of spinal tuberculosis and assess the response of treatment and change in regime of A.T.T. accordingly. **Material and Methods:** In this study, the MRI scans of 49 consecutively managed spinal tuberculosis patients with or without neurological deficit over 1.5 year period were studied cross-sectionally in the Department of Orthopaedics, S. N. Medical College, Agra were analyzed, to determine the pattern of occurrence of various pathological lesions. They were followed up with anti-TB treatment (ATT) according to standard guidelines. **Result:** It was found that spinal tuberculosis was most commonly observed in the 31-50 years age group with a male predominance. The dorsal and the lumbar vertebrae are commonly involved and multiple vertebrae were often affected with kyphotic deformity in 36% cases. Intervertebral disc involvement with pre and paravertebral collections were commonly seen in about 75 % of the cases. Cord oedema was noted in 20% of the cases. Nerve root compression was seen in 63% of cases. Thirty-five patients had no neurological deficits while 14 had paraplegia of varying severity. The canal encroachment was observed in 75% cases. Serial MRI were done during the course of treatment to assess the disease progression. **Conclusion:** This study showed that MRI is the most valuable investigation in the evaluation of spinal tuberculosis. The MRI scan is highly sensitive in the detection of various pathological conditions of spinal tuberculosis and their pattern of occurrence. Serial MRI scans can be used to assess the disease response to treatment.

Key Word: spinal tuberculosis.

*Address for Correspondence:

Dr Saurabh Dwivedi, Department of Orthopaedics, SN Medical College Motikatra 282002, Agra, Uttar Pradesh, INDIA.

Email: dwivedi7500@gmail.com

Received Date: 02/03/2020 Revised Date: 11/04/2020 Accepted Date: 16/06/2020

DOI: <https://doi.org/10.26611/1031512>

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). 

Access this article online

Quick Response Code:	Website: www.medpulse.in
	Accessed Date: 14 July 2020

INTRODUCTION

Tuberculosis is a chronic granulomatous disease caused by infection with *Mycobacterium* species. Spine is the commonest site of skeletal involvement in tuberculosis and constitutes more than 50% of all forms of osteo-articular tuberculosis in India. Annually about 6 million people

become ill with tuberculosis and two million die from disease worldwide.¹ In the developing countries, the disease has an aggressive course, particularly in children and young adults resulting in abscess formation. Consequently, neurologic complications and spinal deformities are frequently observed² Spinal involvement occurs in less than 1% of patients with TB^{3,4} but the increasing frequency of TB in both developed and developing countries has continued to make spinal TB a health problem.^{3,5} Spinal tuberculosis is the most common as well as the most dangerous forms of skeletal TB and accounts for about 50% of all cases of skeletal tuberculosis. Although the dorsolumbar spine seems to be the most common site of the spinal column involvement in spinal tuberculosis, any part of the spine can be affected. Furthermore, the incidence of neurological complications in spinal TB varies from 10% to 44%⁴. Paraplegia may develop in 20 – 25% cases⁵. Tuberculosis of spine is the

most common cause of non-traumatic paraplegia in the most parts of world. Conventional X-rays are useful in diagnosis of spinal tuberculosis but their main disadvantage is that more than 50% of bone has to be destroyed before a lesion can be seen on a plain radiograph, a process which takes nearly six months. Advanced imaging techniques like Magnetic Resonance Imaging (MRI) make the early diagnosis of spinal TB easier and a considerable number of patients with spinal TB are diagnosed earlier and treated before significant neurological deficits develop. However, patients can still present late with considerable spine deformity⁶. MRI because of the lack of ionizing radiation, high contrast resolution, ability to detect marrow infiltration and ease of assessment of extradural disease and status of spinal cord, has become the established optimal imaging technique in the diagnosis of spinal infections and their sequel. However, to date, differentiating tuberculous spondylitis from pyogenic spondylitis is a challenge, in view of varied presentation of the former. MRI has the ability to provide excellent contrast resolution in all planes, making the diagnosis and evaluation of tuberculous spondylitis and para-vertebral extension easier. Serial MRI examinations can be used to assess the response to treatment^{7,8} and are very useful in the management of multilevel infection. Serial MRI scans can also be used to assess the healing of infection.

MATERIALS AND METHODS

A hospital based cross sectional study was evaluated. These cases were from the OPD and IPD from the Department of Orthopedics and the patients who were referred to the Department of Orthopaedics from Department of General Medicine, Pulmonary Medicine and Paediatric Department of S.N. Medical College, Agra on the basis of their clinical presentation over a period of 17 months from November 2017 to May 2019 who were fulfilling inclusion criteria. The inclusion criteria of study was those patients aged 21-60 years of age with or without neurological deficit at spinal level with strong clinical suspicion of spinal tuberculosis. A detailed history along with complete clinical examination and laboratory investigations will be done before the MRI examination. Forty-nine patients with paradiscal spinal TB lesion, with/without paraplegia, aged over twenty years were

enrolled. All patients having classical clinico-radiological diagnosis of spinal tuberculosis or who showed evidence of healing on treatment or with a proven histopathological/bacteriological/cytological diagnosis of TB spine were included in the study. The cases were (A) freshly diagnosed cases of TB spine and (B) patients who had completed treatment successfully and had a pre treatment MRI examination. Trauma patients, patients with metallic implants, patients with pacemaker or cochlear implants, patients with isolated intraspinal granuloma, isolated posterior complex involvement, isolated cranio-vertebral junction TB, isolated sacral tuberculosis or patients without a classical clinico-radiological diagnosis were not included in the study. Neurological assessment was performed by Jain and Sinha scoring system⁶.The radiological diagnosis was made on observations of demineralisation of the vertebra, fuzzy paradiscal margin, reduction or obliteration of disc space, along with one or more of the following findings:- destruction of end plates, wedging of vertebra, obvious kyphotic deformity, paravertebral shadows, anterior scalloping of vertebral body and pre and paravertebral abscess. The aspirate of abscesses (n-9) was sent for Z-N staining, pus culture and PCR. In all prospective patients, MRI was performed pre-treatment and repeated after eight months. Magnetic Resonance Imaging (MRI) examinations were done with 1.5 Tesla Philips Achieva machine using shoulder coil without contrast and the following sequences were taken:-

1. Sagittal and axial T1 weighted (T1 FRFSE),
2. Sagittal and axial T2 weighted (T2 FRFSE),
3. Coronal and sagittal STIR sequences

The following features were assessed by MRI:

Compartment of spine involved: Epidural/ Intradural/ Intramedullary/ Multiple

Epidural involvement assessed for the following

1. Extent of vertebral involvement: body / posterior involvement – signal changes.
2. Wedging or compression.
3. Involvement of disc.
4. Subligamentous extension.
5. Extent of abscess: Epidural / paravertebral / psoas.
6. Spinal cord changes.

Intradural/ intramedullary involvement

Ethical approval was taken from Institution Ethical committee.

RESULTS

Table 1: Age and Sex Distribution

Age Group	Male (29)	Female (20)	Percentage (overall)
21-30	2	1	6
31-40	16	11	55
41-50	7	5	24.5
51-60	4	3	14

Table 2: Regional Distribution of TB Spine

Regional Distribution	No of Cases	Percentage
Cervical	5	10
Thoracic	16	32.5
Lumbar	28	57
Multiple Levels	7	14
Kyphotic Deformity	19	38.5

Table 3: Clinical Profile of Patients with Spinal Tuberculosis

	Fever	Backache	Malaise	Weight Loss
No of cases	25	31	19	9
Percentage	62.5	77	47.5	22.5

Table 4: Extent of Tuberculosis Spine In Various Compartments

Features	No of cases	Percentage
Intervertebral disc involvement	37	75.5
Wedge collapse of body	22	44.8
Complete destruction of vertebra	9	18
Subligamentary extension	21	43
Epidural collection	34	69
Intradural involvement	4	8
Intramedullary involvement	2	4
Pre and Paravertebral involvement	25	52
Discitis in preserved discs	47	95
Nerve root compression	31	63
Canal encroachment	37	75.5
Coed edema	10	20

MR FINDINGS ON HEALING

All cases were put on Anti Tuberculous Therapy (ATT) regimen as per Directly Observed Treatment Short course (DOTS) which consisted of daily dosage with Rifampicin [R] (8-12 mg/kg/day), Isoniazid [H] (4-6mg/kg/day), Pyrazinamide [Z] (20-30 mg/kg/day), Ethambutol [E] (13-17 mg/kg/day) and Streptomycin [S] (20-40 mg/kg/day). The healing of the vertebral lesion was recorded on complete resolution of marrow oedema, replacement of marrow by fat seen as a bright signal on T1WI and T2WI and complete resolution of paravertebral collections. On follow-up MRI seven showed complete resolution of cord oedema and neural deficit. Cord atrophy was observed in three cases. The other some cases developed cord atrophy on completion of treatment. Demineralisation was seen in most of the cases. Myelomalacia, arachnoiditis, syringomyelia, thickened arachnoid complex and meningeal involvement were not found as contrast was not administered for pre-treatment MRI.

DISCUSSION

Tuberculosis has prevailed as a major public health problem especially in our country. Tuberculosis of the spine is clinically important form of extrapulmonary tuberculosis accounting for majority of the

musculoskeletal tuberculosis cases. First described in 1782 by Percival Pott, a British orthopedic surgeon usually occurs due to haematogenous seeding of the vertebra from a distant source. Typically, more than one vertebra is involved and usually affects the anterior aspect of the vertebral body adjacent to the subchondral plate and from there on spreads to involve adjacent intervertebral discs. Further with involvement of bone, wedge collapse and vertebral destruction occurs which results in kyphosis (figure2). Epidural abscess formation (figure2) results in narrowing of the spinal canal diameter with resultant cord compression and neurological deficits. In the present study, we have attempted to depict the various spectrum of presentations of spinal TB with clinical correlation. The regional distribution of vertebra in our study was similar to the findings of DJ Kotzke⁹ and Sajid Ansari.¹⁰ Shanley DJ¹¹ evaluated radiographic manifestations of tubercular spondylitis like intraosseous and paraspinal abscess formation as seen in our study mentioned in Figure 1, paraspinal abscesses in the lumbar region gravitate along the psoas sheath (figure 3) which can extend to the femoral region and cause erosion of overlying skin.^{9,11}

MRI is the gold standard of imaging in tuberculous spondylitis due to its superior soft tissue resolution and multiplanar capability. MRI is very useful to see the

classic pattern of spread of disease starting anteriorly and moving to opposing vertebrae via subligamentous route. As was observed in our study, T1- weighted images usually show hypointense signal within the affected vertebral marrow. In our study, we had two cases showing intramedullary involvement (Figure 4), on MRI tuberculomas appear as low to intermediate signal intensity on T1W images and low signal on T2W images due to caseous necrosis in the tuberculoma, which has high protein content. The extent of spinal cord involvement, nerve root integrity, posterior elements involvement and also response to ATT therapy are best assessed by MRI. It is important to differentiate tuberculous spondylitis from pyogenic spondylitis because proper treatment of the different types can reduce the rate of disability and functional impairment.^{12,13} MRI

has been shown to be accurate in differentiating tuberculous spondylitis from pyogenic spondylitis. If there are presence of well-defined paraspinal abnormal signal, smooth and thin walled abscess, subligamentous spread of three or more vertebral levels and multiple vertebral or entire body involvement it is more suggestive of tuberculous spondylitis than pyogenic spondylitis.¹³ Early recognition and prompt treatment are therefore necessary to minimize residual spinal deformity and permanent neurological deficit Early diagnosis and conservative treatment by ATT has shown favorable results as anti- tuberculous drugs will be able to reach the tuberculous caseous material and cavities in spine. However in patients with severe bone involvement along with cord or root compression, surgical treatment is the only benefitable measure.¹⁴




Figure 1:




Figure 2

Figure 3

Figure 4

Figure 1: MRI of spinal tuberculosis: (a, c-sagittal; b, d-axial; e-coronal) images showing vertebral lesion hypo on T1 and hyper on T2W images, epiphyseal and discal involvement with anterior and paraspinal extension; **Figure 2:** MRI of the thoracic spine in a paraparetic patient demonstrating a pre-vertebral and epidural abscess, bony destruction with kyphotic collapse and the spinal cord draped over an internal gibbus with intrinsic cord signal; **Figure 3:** T2-weighted coronal MR image showing collapse of L1 vertebral body with irregularity of superior end plate of L2 along with bilateral psoas abscesses; **Figure 4:** T2 weighted sagittal plane MRI of the thoraco-lumber spine shows intramedullary peripherally hypointense lesion with central hyperintensity and peripheral edema at T12-L1 vertebrae

CONCLUSION

This study showed that MRI is the most valuable investigation in the evaluation of spinal tuberculosis. The MRI scan is highly sensitive in the detection of various pathological processes of spinal tuberculosis and their pattern of occurrence and also provides information of soft tissue and bone involvement, cord involvement and nerve

root integrity and helps as staging of disease for planning treatment. It is an accurate investigation to differentiate spinal tuberculosis from pyogenic spondylitis and helps in diagnosing spinal TB in early stages and hence prompt treatment minimizes spinal deformity and permanent neurological deficits. Serial MRI scans can also be used

during follow up of patients to monitor the response to treatment.

REFERENCES

1. Ho EKW, Leong JCY. The pediatric spine: principles and practice. In: Weinstein SL, ed. Tuberculosis of the spine, 3rd ed. New York: Raven, 1994:837-49.
2. Sai Kiran NA, Vaishya S, Kale SS, Sharma BS, Mahapatra AK. Surgical results in patients with tuberculosis of the spine and severe lower-extremity motor deficits: a retrospective study of 48 patients. *J Neurosurg Spine*. 2007;6:320-32
3. Bennet JC, Bennet SJ. Infection of Bone and Joint. *Surgery international*, 2006;73, Elsevier Ltd. ISSN 09554572
4. Rezai AR, Lee M, Cooper PR, Errico TJ, Koslow M. Modern management of spinal tuberculosis. *Neurosurgery*. 1995;36:87-97.
5. Turgut M. Spinal tuberculosis (Pott's disease): its clinical presentation, surgical management, and outcome. A survey study on 694 patients. *Neurosurg Rev*. 2001;24:8-13.
6. Boxer DL, Pratt C, Hine AL. Radiological features during and following treatment of spinal tuberculosis. *Br J Radiol* 1992;65(774):476-9.
7. Jallah RD, Kuppasamy I, Mahayuddin AA. Spinal tuberculosis: a five years review of cases at the National Tuberculosis Centre. *Med J Malaya* 1991;46(3):269-73.
8. Lifeso RM, Weaver P, Harder EH. Tuberculous spondylitis in adults. *J Bone Joint Surg*. Dec 1985; 67 A(9):1405-1413.
9. Kotzé DJ, Erasmus LJ. MRI findings in proven Mycobacterium tuberculosis (TB) spondylitis.
10. Sinan TH, Al Ismail M, *et al.* Spinal tuberculosis: CT and MRI features. *Ann Saudi Med*. 2004;24(6):437-41.
11. Shanley DJ. Tuberculosis of the spine: imaging features. *AJR. Am J Roentgenol*. 1995;164:659-64.
12. Arizono T, Oga M, Shiota E, Honda K, Sugioka Y. Differentiation of vertebral osteomyelitis and tuberculous spondylitis by magnetic resonance imaging. *Int Orthop*. 1995;19:319-22.
13. Jung NY, Jee WH, Ha KY *et al.* Discrimination of Tuberculous Spondylitis from Pyogenic Spondylitis on MRI. *AJR*. 2004;182:1405-10.
14. Sinha S, Singh A, Gupta V. Surgical management and outcome of tuberculosis at lantoaxial dislocation: A 15 years experience. *Neurosurgery*. 2003;52(2): 331-9.

Source of Support: None Declared
Conflict of Interest: None Declared

Policy for Articles with Open Access:

Authors who publish with MedPulse International Journal of Orthopedics, (Print ISSN: 2579-0889, Online) (ISSN: 2636-4638) agree to the following terms: Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a Creative Commons Attribution License that allows others to share the work with an acknowledgement of the work's authorship and initial publication in this journal.

Authors are permitted and encouraged to post links to their work online (e.g., in institutional repositories or on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work.