

Role of zinc supplementation in school aged children suffering from common cold: A clinical trial

V K Tandon¹, Gaurav Arya^{2*}

¹Associate Professor, ²Assistant Professor, Department of Paediatrics, Rama Medical College Hospital & Research Centre, Mandhana Kanpur Up, Inida.

Email: arya.drgaurav@gmail.com

Abstract

Background: In developing countries Zinc deficiency is very common among school aged children because of less animal food in diet, and more dietary phytate content. Zinc deficiency causes decreased overall immunity and increases chances of infection. So more studies are needed to see effect of zinc supplement in common cold in children's as it is still questionable. To determine role of supplementation could reduce frequency rate and duration of Common cold during cold season in school aged children living in a low socioeconomic areas of Lakhanpur, Kanpur. **Methods:** We designed a randomized double-blind, placebo-controlled efficacy trial. Subjects were 200 grade 2 primary school children who all completed the trial. Intervention supplementation was zinc sulfate tablets (10 mg elemental) and placebo tablets for case and control groups, respectively. Tablets were taken on a daily basis, 6 days a week, for 5 months (November 2016 to March 2017). **Findings:** Among the zinc-supplemented group common cold incidence of 1.37 ± 0.86 episodes per child during the study period was recorded in comparison to 3.15 ± 0.55 cold episodes per child among the placebo group ($P < 0.001$). Mean overall missing days from school was 0.55 ± 1.09 days and 1.35 ± 1.79 days for zinc-supplemented and placebo groups, respectively. The need for administration of antibiotics for bacterial infections (pharyngitis, acute otitis media, sinusitis, pneumonia) were 20 and 47 courses for zinc-supplemented and placebo groups, respectively ($P < 0.01$). **Conclusion:** This study showed that zinc supplementation has a beneficial impact on the occurrence of common cold **Key Words:** Zinc Supplementation; common cold; upper respiratory Infection; children.

*Address for Correspondence:

Dr Gaurav Arya, Assistant Professor Department of Paediatrics, Rama Medical College Hospital & Research Centre, Mandhana Kanpur Uttar Pradesh, INDIA.

Email: arya.drgaurav@gmail.com

Received Date: 20/11/2019 Revised Date: 19/12/2019 Accepted Date: 11/01/2020

DOI: <https://doi.org/10.26611/10141421>

Access this article online

Quick Response Code:



Website:

www.medpulse.in

Accessed Date:
06 May 2020

INTRODUCTION

Zinc is an essential micronutrient for human growth, development, and immune function. Mild to moderate zinc deficiency can be best detected through a positive response to supplementation trials. Zinc supplementation has been

shown to have a positive effect on the incidence of diarrhea, and pneumonia¹. Upper respiratory tract viral infections are one of the most common reasons for physician visits. School aged children may experience 6 to 8 colds per year². In children, this illness is also more extensive than in adults and usually requires medical attention. Statistics indicate that more than 80% of common colds requiring medical attention affect children and adolescents³. Zinc salts have been found to inhibit rhinovirus replication *in vitro*, possibly by interfering with rhinovirus protein cleavage⁴. It has also been suggested that cold symptoms, sneezing and nasal congestion, might be reduced by elevations in intranasal zinc salts, producing a chemical clamp, or may be due to correction of a subclinical zinc deficiency^{5,6}. Several controlled trials of treatment of common cold with zinc have been published; however, there are only a few community-based

longitudinal studies that have been conducted with zinc supplementation for prevention of common cold during cold season⁷. To investigate the effect of zinc supplementation on the prevention of occurrence and the need for antibiotic administration, we performed an intervention trial among school aged children in lakhanpur, kanpur.

Subjects and Methods

The study was a randomized, double-blind, placebo-controlled, community-based intervention trial conducted between November 2016 and March 2017 in a low socioeconomic areas of lakhanpur, Kanpur. The project was approved by the scientific advisory and ethical committees of Rama University of Medical Sciences Kanpur. Written informed consent forms were signed by the parents before including the children in the study. For ethical reasons and limitation of cost we did not measure plasma Zinc levels in this study. A total of 200 children (aged 78 to 120 months) were randomly assigned to daily (6 days in week) supplementation with 10 mg elemental zinc as a tablet (n=100, 50 males, and 50 females), or placebo (n=100, 50 males, and 50 females). For each child

that was enrolled in the study, a standardized questionnaire was completed to obtain family details including basic demographic and socioepidemiologic data. The subjects were free of chronic diseases, such as sickle cell disease, or protein-energy malnutrition. All participants were observed on a daily basis for any medical sign and symptom especially cold symptoms. This daily surveillance for the detection of any sign and symptom of common cold and other possible diseases was conducted by trained health workers. These workers were also responsible for proper administration of the zinc or placebo tablets. Regular field visits (once a week) for additional data collection and supervision of the health workers were made by our physician. Common cold was defined as the presence of at least two of the following symptoms: cough, headache, hoarseness, muscle ache, nasal drainage, nasal congestion, scratchy throat, sore throat, sneezing, and fever. Data were analyzed using statistical software packages SPSS 11.5 (SPSS, Inc, Chicago, IL) and Epi Info (Center for Disease Control and Prevention, Atlanta, GA) using the χ^2 test, *t*-test and ANOVA, *P* values less than 0.05 was considered statistically significant.

Findings

Table 1 summarizes the demographic characteristics of participants. The average occurrence of common cold was 1.73 ± 0.86 in zinc recipients and 3.15 ± 0.55 in placebo recipients ($P < 0.001$)

Table 1: Demographic findings of 200 students treated with zinc sulfate or Placebo

Variable	Zinc group	Placebo group	P value
Age (month)	93.7(7.38)	93.1(7.35)	0.09
Mean (SD)			
Sex Male	50	50	
Female	50	50	
Number of family members	4.7(2.2)	4.5(2.1)	0.7
Smoker parent	18	19	0.1

* SD: Standard Deviation

(Table 2). Missing days from school during 5 months of study were 0.55 ± 1.09 days (per student) and 1.35 ± 1.79 days (per student) in zinc-supplemented and control groups, respectively ($P < 0.001$). In the zinc supplemented group, three participants complained from mild gastrointestinal discomfort which was resolved within few days and there was no need for their exclusion from the study.

DISCUSSION

Millions of people throughout the world may have inadequate levels of zinc in their diet due to limited access to zinc-rich foods (animal products, oysters and shellfish) and the abundance of zinc inhibitors such as phytate, common in plant-based diets⁸. Zinc is a key component of the cell architecture and is required for the production of over 200 enzymes including phosphatase,

metalloproteinases, oxidoreductase, and transferase which are involved in protein synthesis, nucleic acid metabolism, and immune function¹⁸. Zinc deficiency is common in young children in the developing countries and is associated with reduced immuno-competence and increased rates of serious infectious diseases⁹. Several studies have shown that zinc supplementation has a positive influence on linear, motor development and weight gain¹⁹. Community-based studies conducted among children of different age groups have shown the beneficial impact of zinc supplementation in the form of reduced diarrhea episodes in children with zinc deficiency, since this supplementation may lead to accelerated regeneration of mucosa, increased levels of brush border enzymes, enhanced cellular immunity, and higher levels of secretory antibodies¹⁰.

Table 2: Median common cold occurrence need for antibiotic missing school Among 200 Students Treated with Zinc sulfate or Placebo

Variable	Zinc group	Placebo group	P value
Average common cold occurrence(SD)	1.7(0.86)	3.1(0.55)	<0.001
Max			
Min	6	8	
	0	0	
Need for antibiotic administration	20	47	<0.001
Days missing school	0.55+1.09	1.35+1.79	<0.001

* SD: Standard Deviation

We found that supplementation with zinc was associated with a decrease in the average occurrence of common cold during cold months of year, among children living in a low socioeconomic area of lakhanpur, Kanpur. We also found that supplementation with zinc was associated with a decrease in the average duration and severity of common cold. Prasad and co-workers showed a beneficial effect of zinc lozenges for reduction of duration and severity of cold symptoms⁷. They believed improvement in cold symptoms was related to the antioxidant anti-inflammatory effect of zinc¹⁹. However, previous trials failed to show a beneficial effect of zinc for treatment of common cold, perhaps because inadequate doses or inappropriate formulations of zinc were used, resulting in lack of bioavailable zinc^{11,12}. Three other double-blind placebo-controlled trials evaluated the prophylactic and therapeutic efficacy of zinc gluconate glycine lozenges. In the first study involving 57 volunteers, administration of zinc gluconate lozenge (23 mg every 2 hours for 4.5 days) started one day before inoculation with human rhinovirus, reduced the total mean clinical score to 5.7 from 8.2 with placebo^{13,14}. In another study 69 subjects were inoculated with human rhinovirus, and the 12 who experienced cold symptom were randomly allocated to receive either zinc gluconate lozenges or matched placebo every 2 hours for 6 days while awake. Once again zinc treatment significantly reduced clinical scores compared to placebo¹⁵. Retrospective chart analysis study by McElroy and co-workers provides strong support for the beneficial effects of zinc in school-aged subjects with common cold. They concluded that treatment with zinc can reduce duration of cold signs and symptoms and the need for antibiotics, and prophylaxis may decrease the incidence of colds¹⁶. Hulisz showed that zinc administration within 24 hours of the onset of common cold may reduce the duration and severity of symptoms of common cold¹⁷. According to American Family Physician guideline zinc is not recommended for treatment of common cold because of these inconsistent study results²⁰. In our study only 20 participants in zinc supplementation group required antibiotics for treatment of upper respiratory tract infection compared to 47 individuals that needed antibiotic therapy in placebo group. Recently, Kurugol and coworkers carried out a prospective study to determine the prophylactic and therapeutic effectiveness of zinc sulphate for the common

cold in children. The results of the study showed that the use of zinc sulphate significantly reduced the risk of colds and cold related school absences³.

CONCLUSION

In conclusion we showed that supplementation with zinc was associated with significant decrease of the frequency of colds, and reduction in the duration and severity of the symptoms of common cold. In addition, the need for use or misuse of antibiotics was reduced. Fortification of more commonly used foods such as bread with zinc salts may be beneficial. However, further investigation is required.

Acknowledgment

This study was supported by Rama medical university Mandhana and primary school Lakhanpur Kanpur. Participation of the children and their parent is also acknowledged.

REFERENCES

1. Walker CF, Black RE. Zinc and the risk for infectious disease. *Ann Rev Nutr.* 2004;24: 255- 75.
2. Dingle JH, Badger GF, Jordan WS. Illness in the home: study of 25000 illnesses in a group of Cleveland families. Cleveland, Press of WesternReserve University, 1964.
3. Kurugöl Z, Akilli M, Bayram N, Koturoglu G. The prophylactic and therapeutic effectiveness of zinc sulphate on common cold in children. *Acta Paediatr.* 2006;95(10): 1175-81.
4. Korant BD, Butterworth BE. Inhibition by zinc of rhinovirus protein cleavage: interaction of zinc with capsid polypeptides. *J Virol.* 1976;18(1):298-306.
5. Novik SG, Godfrey JC, Godfery NJ, Wilder HR. How does zinc modify the common cold? *Med Hypoth.* 1996;46(3):295-302.
6. Prasad AS. Zinc: The Biology and Therapeutics of an Ion. *Ann Internal Med.* 1996;125(2):142-4.
7. Prasad AS, Fitzgerald JT, Bao B, *et al...* Duration of symptoms and plasma cytokine levels in patients with the common cold treated with zinc acetate. *Ann Int Med.* 2000; 133(4):245-52.
8. Sandstead HH. Zinc deficiency a public health problem? *Amer J Dis Child.* 1991; 145(8):853-9.
9. Sur D, Gupta DN, Mondal SK, *et al...* Impact of zinc supplementation on diarrheal morbidity and growth pattern of low birth weight infants in Kolkata, India: A randomized double-blind, placebo controlled, community-based study. *Pediatrics.* 2003; 112(6 pt 1):1327-32.

10. Folwaczny C. Zinc and diarrhea in infants. *Trace Elem Med Biol.* 1997;11(2):116-22.
11. Douglas RM, Miles HB, Moore BW, *et al.* Failure of effervescent zinc acetate lozenges to alter the course of upper respiratory tract infection of Australian adults. *Antimicrob Agents Chemother.* 1987;31(8):1263-5.
12. Maknin ML, Piedmonte M, Calendine C, *et al.* Zinc gluconate lozenges for treating common cold in children. *JAMA* 1998;279(24):1962-7.
13. AL-Nakib W, Higgins PG, Barrow I, *et al.* Prophylaxis and treatment of rhinovirus cold with zinc gluconate lozenges. *Antimicrob Chemother.* 1987;20(6):893-901.
14. AL-Nakib W, Higgins PG, Barrow I, *et al.* Prophylaxis and treatment of rhinovirus cold with zinc gluconate lozenges. *Am J Ther.* 2003; 10(3):233-4.
15. McELroy BH, Miller SP. Effectiveness of zinc gluconate glycine lozenges (Cold- Eeze) against the common cold in school- aged subjects: a retrospective chart review. *AM J Ther.* 2002;9(6):472-5.
16. Hulisz D. Efficacy of zinc against common cold viruses: an overview. *J Am Pharm Assoc.* 2004;44(5):594-603.
17. Diaz-Gomez NM, Domenech E, Barroso F, *et al.* The effect of zinc supplementation on linear growth, body composition and growth factors in preterm infants. *Pediatrics.* 2003; 111(5 pt1): 1002-9.
18. Lira PIC, Ashworth A, Morris SS. Effect of zinc supplementation on the morbidity, immune function, and growth of low-birth-weight, fullterm infants in northeast Brazil. *Am J Clin Nutr.* 1998; 68(2 suppl): 418S-424S.
19. Prasad AS, Beck FW, Bao B, *et al.* Duration and severity of symptoms and levels of plasma interleukin-1 receptor antagonist, soluble tumor necrosis factor receptor, and adhesion molecules in patients with common cold treated with zinc acetate. *J Infec Dis.* 2008;197(6):795-802.
20. Simasek M, Blandino DA. Treatment of common cold. *Am Fam Physician.* 2007;75(4): 515-20.

Source of Support: None Declared
Conflict of Interest: None Declared

Policy for Articles with Open Access:

Authors who publish with MedPulse International Journal of Pediatrics (Print ISSN: 2579-0897) (Online ISSN: 2636-4662) agree to the following terms: Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a Creative Commons Attribution License that allows others to share the work with an acknowledgement of the work's authorship and initial publication in this journal.

Authors are permitted and encouraged to post links to their work online (e.g., in institutional repositories or on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work.

