

Developmental assessment of children with neonatal hypoglycemia

Rasika Bharaswadkar^{1*}, Sharad Agarkhedkar², Shradha Salunkhe³, Manoj Patil⁴, Vineeta Pande⁵, Shiji Chalipat⁶, Vishwanath Kulkarni⁷, Sandeep Patil⁸

^{1,7}Assistant Professor, ²Professor & HOD, ^{3,4,6}Associate Professor, ⁵Professor, ⁸Pediatric Neurologist, Department of Pediatrics, Dr. D.Y. Patil Medical College, Hospital And Research Center, Dr. D. Y. Patil Vidyapeeth, Pimpri, Pune 411018, Maharashtra, INDIA.

Email: dr.rasika80@gmail.com

Abstract

Background: Neonatal hypoglycemia is a frequently encountered metabolic derangement confronted in neonates leading to developmental delay in later life. Early identification of delay can help the treating paediatrician to start early stimulation and improve the outcome of these children. Various developmental screening tests are available but these require expertise administration and interpretation. **Aims:** Compare DDST II with TDSC for screening for developmental delay and LEST for screening of language delay in children with neonatal hypoglycemia. **Methods:** We conducted a Descriptive Hospital-based cross-sectional study after Institutional ethics committee approval and written informed consent. Children with history of neonatal hypoglycemia were identified and their details were collected in a structured proforma. All these children were assessed for developmental delay by DDST II, TDSC and LEST scale by different individuals. Statistical analysis was done by using SPSS software. **Results:** Total of 81 children were enrolled in the age group of 6 months to 6 years. On comparing TDSC to DDST II, sensitivity of the TDSC is 93.18% and specificity is 100%. Kappa value is 0.92 (0.70- 1.14). While LEST has sensitivity is 88.64 % specificity is 97.3 % and Kappa value is 0.852 (1.071- 0.6367) as compared to the Language domain of DDST II. **Conclusions:** TDSC and LEST are simple scales with good sensitivity and specificity. Nurse, receptionist or peripheral health workers can be taught to use these scales with minimum training to augment early identification and early referral of developmental delay.

Key Words: developmental delay, language delay, Specificity, sensitivity.

*Address for Correspondence:

Dr Rasika Bharaswadkar, Assistant Professor, Department of Pediatrics, Dr D.Y. Patil Medical College, Dr. D. Y. Patil Vidyapeeth, Sant Tukaram Nagar, Pimpri, Pune 411018, Maharashtra. INDIA.

Email: dr.rasika80@gmail.com

Received Date: 01/10/2020 Revised Date: 11/11/2020 Accepted Date: 07/12/2020

DOI: <https://doi.org/10.26611/10141711>

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). 

Access this article online

Quick Response Code:	Website: www.medpulse.in
	Accessed Date: 02 January 2021

INTRODUCTION

Neonatal hypoglycemia is one of the frequently encountered metabolic derangements confronted in neonates. Hypoglycemia occurs in 1.3 – 4.4 per 1000 full-term newborns and 15–55 per 1000 preterm newborns¹.

Hypoglycemia had a deleterious effect on the brain. Neonatal hypoglycemia leads to various neurodevelopmental disabilities in children like microcephaly, cerebral palsy, behavioural disorders, seizures, and visual disturbances and developmental delays. Early identification of these delays can help the treating paediatrician to start early stimulation and improve the outcome of these children. The various screening tests used for developmental assessment are available like Denver developmental screening test (DDST II) and Bayley developmental screening test. The DDST II is a valid scale with a strong relationship between classification on the DDST and scores on the Stanford-Binet intelligence scales and the Bayley infant scales². DDST-II classifies the assessment of child's development into 4 areas personal-social (25 items), fine motor-adaptive (29 items), language (39 items), and gross motor

(32 items)³. These tools require more expertise in administering the test. Screening tools like Trivandrum Developmental Screening Chart (TDSC) for children of 0-6 years and Language evaluation Scale Trivandrum (LEST) 0-6 years are simple tools which can be administered by the nursing staff. This study was undertaken to compare DDST II with TDSC for screening for developmental delay and LEST for screening of language delay.

MATERIALS AND METHODS

Aim: Compare DDST II with TDSC for screening for developmental delay and LEST for screening of language delay in children with neonatal hypoglycemia.

Objective:

1. To identify children with neonatal hypoglycemia and classify them into symptomatic and asymptomatic neonatal hypoglycemia.
2. To assess the various domains of DDST II and compare them among the two groups.
3. To study the sensitivity and specificity of TDSC for screening for developmental delay in comparison with DDST II in children with neonatal hypoglycemia.
4. To study the sensitivity and specificity of LEST for screening for language delay in comparison with DDST II in children with neonatal hypoglycemia.

Type of study: Descriptive Hospital-based cross-sectional study.

Inclusion criteria:

1. Children with history of neonatal hypoglycemia.
2. Age group 6 months to 6 years

Exclusion criteria:

Children with severe congenital anomalies and syndromic children were excluded from the study.

Place of study:

Dr. D. Y. Patil Medical College and Hospital.

METHODOLOGY

All children coming to the OPD or IPD of Dr. D.Y. Patil Medical College and Hospital Pimpri, Pune from 2016 to 2018 were our study population. They were screened for history of neonatal hypoglycemia. All subjects with documented evidence of neonatal hypoglycemia were enrolled in the study after parents voluntarily signed the written informed consent. Details of their sociodemographic profile and perinatal history were documented in a structured proforma. Subjects were classified into Symptomatic and Asymptomatic hypoglycemia depending on the presence or absence of

symptoms of hypoglycemia during neonatal period. Details of the presenting complains were noted. Structured neurological examination was done in all patients.. All patients underwent detail developmental screening by using DDST II, TDSC and LEST charts at the same setting. A line is drawn at the chronological age of the child on the chart and developmental millstones assessed. If the child fails any item on the left side of the line it is labelled as Delay on DDST II and TDSC^{4,5}. For LEST 1 item is taken as suspect and 2 items as delay^{6,7}. Assessment by DDST II, TDSC and LEST was done by independent individuals who were trained on how to administer the scale to avoid bias. All these findings were recorded in the case sheet and later entered into excel. Detail statistical analysis was done using SPSS software.

OBSERVATIONS AND RESULTS

We included 81 children in the study who had definite history of neonatal hypoglycemia. We had 62(76.54%) males out of 81. The maximum number ie: 31 (38.3 %) belong to the 6 to 12 months age group, followed by 24 (29.6%), 7(8.6 %) and 6 (7.4 %) in age group of 13 – 24 months, 25- 36, 37- 48, 49 – 60 and 61- 72 months respectively. Out of the total 81 patients 48 (59.3%) patients had symptomatic hypoglycemia and 33 (40.7 %) had asymptomatic hypoglycemia. In our study total 44 (54.3%) children had developmental delay by using DDST II as screening tool. Statistically significant delay was seen in the number of children with symptomatic hypoglycemia {32 (72.7%)} as compared to asymptomatic hypoglycemia children. Table 2 shows distribution of various domains of DDST II in detail among children with symptomatic and asymptomatic hypoglycemia. On using TDSC, developmental delay was found in 41 children. Thus 3 children were picked up in addition when we applied DDST II. Details are shown in table 3. Language domain was again assessed by LEST which picked up 37 (45.7%) children with language delay and 3 children were suspect for language delay. DDST II language domain had picked up 40 children with speech delay which matches well with LEST (Table 4). In table 5, out of the 50.6 % cases who had delay on TDSC also had delay on DENVER II. There were only 3 children who were normal on TDSC and were found to have delay on DDST II. Sensitivity of the TDSC is 93.18% and specificity is 100%. Kappa value is 0.92 (0.70- 1.14). This indicates a strong agreement between TDSC and DENVER II. Table 6 shows 39 (97.5 %) children had language delay on LEST. Sensitivity is 88.64 % specificity is 97.3 % and Kappa value is 0.852 (1.071- 0.6367). This indicates a strong agreement between LEST and DENVER II for classification of patients into language delay and normal.

Table 1: Distribution of symptomatic and asymptomatic neonatal hypoglycemia cases according to Denver developmental screening test II (DDST II)

Denver II	SYMPTOMATIC	ASYMPTOMATIC	TOTAL	P Value
DELAY	32 72.7%	12 27.3%	44 (54.3%)	0.0075
NO DELAY	16 43.2%	21 56.8%	37 (45.7%)	
TOTAL	48 (59.3%)	33 (40.7%)	81 (100%)	

Table 1 shows that total 44 children had delay according to DDST II of which 32 (72.7%) children had symptomatic hypoglycemia. There was statistically significant difference between the two groups.

Table 2: Distribution of symptomatic and asymptomatic neonatal hypoglycemia cases according to various domains of DDST II

DENVER II	Symptomatic		Asymptomatic		Total		P Value
	No	%	No	%	No	%	
Gross motor Delay	31	73.8	11	26.2	42	51.9	0.0060
Fine Motor Delay	30	75.0	10	25.0	40	49.4	0.0047
Language Delay	29	72.5	11	27.5	40	49.4	0.0173
Personal Social Delay	30	75.0	10	25.0	40	49.4	0.0047

Table 2 shows distribution of various domains assessed by DDST II among symptomatic and asymptomatic hypoglycemia. Symptomatic hypoglycemia children had statistically significant delay in all the domain's when compared to asymptomatic hypoglycemia children.

Table 3: Distribution of symptomatic and asymptomatic neonatal hypoglycemia cases according to Trivandrum developmental assessment scale (TDSC).

TDSC	SYMPTOMATIC	ASYMPTOMATIC	TOTAL	P Value
DELAY	30 73.2%	11 26.8	41 (50.6%)	0.0104
NO DELAY	18 45.0%	22 55.0%	40 (49.4%)	
TOTAL	48 (59.3%)	33 (40.7%)	81 (100%)	

The table no 3 shows that total 41 children had delay on TDSC of which 30 (73.2 %) children belonged to the symptomatic hypoglycemia. This difference was statistically significant.

Table 4: Distribution of symptomatic and asymptomatic neonatal hypoglycemia cases according to Language assessment scale Trivandrum (LEST)

LEST	SYMPTOMATIC	ASYMPTOMATIC	TOTAL	P value
DELAY	26 70.3%	11 29.7%	37 (45.7%)	0.1502
NO DELAY	21 51.2%	20 48.8%	41 (50.6%)	
SUSPECT	1 33.3% T	2 66.7%	3 (3.7%)	
TOTAL	48 (59.3%)	33 (40.7%)	81 (100%)	

In table 4 we see that 37 (45.7%) of total had language delay. 26 (70.3 %) children among symptomatic group had delay on LEST while 3 children were suspect for delay. The difference between the two groups was not statistically significant.

Table 5: Comparison of neonatal hypoglycemia cases according to DDST II and TDSC

TDSC Outcome	DENVER II outcome			P Value
	Delay	Normal	Total	
Delay	41 100%	0 0.0%	41(50.6%)	0.0001
Normal	3 7.5%	37 92.5%	40 (49.4%)	
Total	44 (54.3%)	37 (45.7%)	81 (100%)	

As indicated in table 5, out of the 50.6 % cases who had delay on TDSC also had delay on DENVER II. There were only 3 children who were normal on TDSC and were found to have delay on DDST II. Sensitivity of the TDSC is 93.18% and specificity is 100%. Kappa value is 0.92 (0.70- 1.14). This indicates a strong agreement between TDSC and DENVER II

Table 6: Distribution of neonatal hypoglycemia cases according to DDST II and LEST

LEST Outcome	DENVER II outcome			P Value
	Delay	Normal	Total	
Delay	39 97.5%	1 2.5%	41 (50.6%)	0.0001
Normal	5 12.2%	36 87.8%	40 (49.4%)	
Total	44 (54.3%)	37 (45.7%)	81 (100%)	

Table 6 shows 39 (97.5 %) children had language delay on LEST. On applying DDST II an additional 5 children were picked up which increased the count to 44 (54.3%). Sensitivity is 88.64 % specificity is 97.3 % and Kappa value is 0.852 (1.071- 0.6367). This indicates a strong agreement between LEST and DENVER II for classification of patients into language delay and normal.

DISCUSSION

In our sample of 81 children we had dominance of male sex {62(76.54%)}, which was similar to study done by Singh *et al.*⁸. S Thirumalaikumarasamy *et al.* found a female preponderance seen in their study⁹. In our study forty four (54.3 %) children had delay on DDST- II, of which 32 (72.7%) were symptomatic (P = 0.0075). Singh *et al.* found that 8 (1.9%) developed symptomatic hypoglycemia out of 107 babies.⁸ Meiri *et al.* found that hypoglycemia was symptomatic in four infants, all of whom were below the fifth percentile for BW¹⁰. On comparing individual domains of DDST II which includes the Gross Motor, Fine Motor, Language, and Personal social domain, all domains are statistically significantly affected in children with symptomatic hypoglycemia. Melana *et al.* in their prospective study of 39 neonates found that the prevalence of abnormal neurodevelopmental outcome in children with neonatal hypoglycemia by DDST 2 method was 71.79% [n=28] and 66.6% [n=26] at 3 and 6 months respectively¹¹. TDSC and LEST were the other scales used to assess development which showed a similar correlation. On comparing the assessment by DDST II with TDSC we found that TDSC has a sensitivity of 93.18% and specificity of 100% over DDST- II. % Kappa value is 0.92 (0.70 - 1.14). This indicates a strong agreement between TDSC and DENVER II. Also, on comparing DDST II with LEST, we found a sensitivity of 88.64 % and specificity of 97.3 % and Kappa of 0.852 (1.071- 0.6367). This indicates a strong agreement between LEST and DENVER II for the classification of patients into Language delay and normal. Nair *et al.* conducted a study on “Development and Validation of Trivandrum Development Screening Chart for Children Aged 0-6 years” using DDST as the reference standard⁵. On delay in one item on TDSC (0-6 y) being considered as ‘TDSC delay’ (test positive), the sensitivity of TDSC (0-6 y) was found to be 84.62 % (95 % CI:

71.92–93.12) and specificity was 90.8 % (95 % CI: 88.97–92.43). The Negative Predictive Value of 99.23 % (95 % CI: 98.48– 99.67) and LR (negative) of 0.17(95 % CI: 0.09–0.32). Nair *et al.* also validated LEST against Receptive Expressive Emergent Language Scale (REELS) for 0-3 years¹² and Extended REELS for 3-6 years age group⁶. The LEST 0-3 screening tool showed a sensitivity of 84.4%, specificity of 80.3%, Positive Predictive Value (PPV) of 91.5%, Negative Predictive Value (NPV) of 67.1% and accuracy (83.2%) against the reference standard REELS. For LEST 3-6 years scale showed a sensitivity of (81%, 47%); specificity (68%,94%), PPV (12%, 31%); NPV (98%, 97%) and accuracy (68.5%, 92%), respectively. Kishore *et al.* in their study “To identify clinical utility of TDSC in screening of developmental delay in children (0-3 yrs.) as compared to DDST” concluded a sensitivity of 57.4% and specificity of 100% for TDSC as against DDST for screening developmental delay. We did not find any study comparing DDST II and LEST for language evaluation in the literature that we reviewed till date¹³. “Development of High-Risk Newborns – A Follow-up Study from Birth to One Year” by Elenjickal *et al.* found the sensitivity of 57.4% and specificity of 100% for TDSC as against DDST for screening developmental delay¹⁴. Ryu and Sim conducted a study on “The validity and reliability of DDST II and Bayley III in children with language development delay”. They proposed that DDST II is a useful screening test to identify infants with delayed language development¹⁵. Shahshahani *et al.* have validated a Persian version of the DDST II for use in Iranian children¹⁶.

CONCLUSION

Developmental screening is very important specially among the high-risk group children. It gives an idea to the care giver about the domain’s affected and a rough idea of

the severity of delay. This helps them work with focus and can reinforce therapies to promote further development. DDST II even though simpler is time consuming and requires adequate training and experience to administer. TDSC and LEST are quick and simpler tests. Thus, we conclude that TDSC and LEST are simple scales with good sensitivity and specificity. These are simpler to use specially in busy OPDs and can be administered by the receptionist or nurse for primary screening of developmental delay in children.

Limitation:

We have not used this scale in the community level population. Targeting to a high-risk group may have some fallacies in the results.

REFERENCES

1. Barbosa ADM, Júnior IF, Lima GM de. Neonatal Hypoglycemia. *Sel Top Neonatal Care* [Internet]. 2017 Dec 20 [cited 2020 Nov 30]; Available from: <https://www.intechopen.com/books/selected-topics-in-neonatal-care/neonatal-hypoglycemia>
2. Salvia J, Ysseldyke JE. Assessment: In *Special and Inclusive Education*. 9th edition. Boston: Houghton Mifflin College Div; 2003. 728 p.
3. Frankenburg WK, Dodds J, Archer P, Shapiro H, Bresnick B. The Denver II: a major revision and restandardization of the Denver Developmental Screening Test. *Pediatrics*. 1992 Jan;89(1):91–7.
4. Nair MKC, Princly P, Leena ML, Swapna S, Kumari I L, Preethi R, et al.. CDC Kerala 17: Early Detection of Developmental Delay / Disability Among Children Below 3 y in Kerala - A Cross Sectional Survey. *Indian J Pediatr*. 2014 Dec;81(S2):156–60.
5. Nair MKC, Nair GSH, George B, Suma N, Neethu C, Leena ML, et al.. Development and Validation of Trivandrum Development Screening Chart for Children Aged 0-6 years [TDSC (0-6)]. *Indian J Pediatr*. 2013 Nov;80(S2):248–55.
6. Nair MKC, Harikumar GSN, George B, Mini AO. Language Evaluation Scale Trivandrum (LEST 3-6 years) Development and Validation. *Indian Pediatr*. 2016 Mar;53(3):257–8.
7. Nair MKC, Harikumar Nair GS, Mini AO, Indulekha S, Letha S, Russell PS. Development and validation of Language Evaluation Scale Trivandrum for children aged 0–3 years — LEST (0–3). *Indian Pediatr*. 2013 May;50(5):463–7.
8. Singh M, Singhal PK, Paul VK, Deorari AK, Sundaram KR, Ghorpade MD, et al.. Neurodevelopmental outcome of asymptomatic and symptomatic babies with neonatal hypoglycaemia. *Indian J Med Res*. 1991 Feb;94:6–10.
9. Thirumalaikumarasamy S, Ramalingam E, Moorthi MMS, Nadesan B. The incidence of asymptomatic hypoglycemia in term newborn babies weighing more than two kilograms. *Int J Contemp Pediatr*. 2017 Jun 21;4(4):1267–73.
10. Mejri A, Dorval VG, Nuyt AM, Carceller A. Hypoglycemia in term newborns with a birth weight below the 10th percentile. *Paediatr Child Health*. 2010;15(5):271–5.
11. Melana N, Ahmed N, Soni RK, Goyal M. Neurodevelopmental Outcome in Neonates with Hypoglycaemia and Associated Risk Factors: A Follow up Study. *J Pregnancy Child Health*. 2017 May 17;4(3):1–6.
12. Nair MKC, Mini AO, Bhaskaran D, Harikumar Nair GS, George B, Leena ML, et al.. CDC Kerala 6: Validation of Language Evaluation Scale Trivandrum (0–3 y) Against Receptive Expressive Emergent Language Scale in a Developmental Evaluation Clinic Population. *Indian J Pediatr*. 2014 Dec;81(S2):99–101.
13. K.1 KK, Sharma2 P, Chavan3 DV, K.4 MB, E.5 A. KEYWORDS Developmental Delay, Screening Tools, TDSC- Trivandrum Development Screening Chart, DDST- Denver Development Screening Chart. IDENTIFY Clin Util TDSC Screen Dev DELAY Child 0-3 Yrs Comp DDST [Internet]. 2019 Jun 8 [cited 2020 Nov 28];(98635). Available from: https://jebmh.com/latest_articles/98635
14. Elenjickal M, Thomas K, Sushamabai S, Ahamed S. Development of High-Risk Newborns - A Follow-up Study from Birth to One Year. *Indian Pediatr*. 2009 Feb 1;46:342–5.
15. Ryu SH, Sim Y-J. The validity and reliability of DDST II and Bayley III in children with language development delay. *Neurol Asia*. 2019;7.
16. Shahshahani S, Vameghi R, Azari N, Sajedi F, Kazemnejad A. Validity and Reliability Determination of Denver Developmental Screening Test-II in 0-6 Year-Olds in Tehran. *Iran J Pediatr*. 2010 Sep;20(3):313–22.

Source of Support: None Declared
 Conflict of Interest: None Declared